

# SPDE LIMITS FOR THE AGE STRUCTURE OF A POPULATION

JIE YEN FAN, KAIS HAMZA, PETER JAGERS, AND FIMA C. KLEBANER

**ABSTRACT.** We consider a family of general branching processes with reproduction parameters depending on the age of the individual as well as the population age structure and a parameter  $K$ , which may represent the carrying capacity. These processes are Markovian in the age structure. In a previous paper [7] the law of large numbers as  $K \rightarrow \infty$  was derived. Here we prove the Central Limit Theorem, namely the weak convergence of the properly scaled processes in an appropriate Skorokhod space. We also show that the limit is driven by an SPDE.

## 1. INTRODUCTION

A branching process is used to model a system of particles where each particle has a random lifespan and gives birth to a random number of offspring at some point during lifetime or at death. Classical frameworks of branching process include the Galton-Watson process in discrete time and the Bellman-Harris branching process in continuous time. In the Bellman-Harris framework, each particle lives for a random length of time and reproduces, independently of each other and with the same law, at death a random number of offspring. In this paper, we consider a much more general framework introduced by Jagers and Klebaner ([12], [13]).

Consider a population of size  $z$  with ages  $(a_1, a_2, \dots, a_z)$ . This age structure can be represented by the measure  $A = \sum_{i=1}^z \delta_{a_i}$  on  $\mathcal{B}$ , the Borel  $\sigma$ -field of  $\mathbb{R}_+$ , where  $\delta_a$  denotes the Dirac measure at  $a$ . In particular, for a measurable set  $B$ ,  $A(B)$  represents the number of individuals with ages in  $B$ . While the size of the population at time  $t$  in the Bellman-Harris process is not Markov, the measure-valued process of ages is. The Markov property remains even when the life span and reproduction of individuals are allowed to depend on the whole population.

We allow for reproduction and death to depend on the age, the age structure of the population, as well as some parameter  $K$ . The latter could play the role of the carrying capacity of the habitat. We allow for multiple offspring during life and at death to have a rather general model. We are interested in the approximations when  $K$  is large. Remark that our CLT yields new results even in the classical case of constant parameters.

Similar questions have been answered in [17] and [18] under the Galton-Watson setting, where the reproduction of each particle depends on the carrying capacity, but is otherwise independent and identically distributed conditionally on the carrying capacity and the size of the population. Oelschläger [24] also answered a similar question in the context of birth-death processes, deriving a LLN and a CLT for the empirical processes of age-structured populations as the size of population tends to infinity.

Tran [25] (also [26] and [6]) obtained a LLN and a CLT for a population model structured by traits and ages (not just the physical age). He generalises the standard

model by including the possibility of trait mutations and interactions (through a kernel) among individuals, while keeping the dependence of the reproduction on just the state (traits and ages) of that individual. In contrast, we allow the births and deaths to depend on the age structure of the whole population. Kaspi and Ramanan ([15] and [16]) obtained LLN and CLT for measure-valued queuing processes, which inspired this paper.

Convergence of measure-valued processes has been studied in various settings over the last decades. This has been done also in the context of population or particle systems, either giving results of the type of LLN only (e.g. [4], [20], [21]), or together with CLT (e.g. [3], [19], [24], [25]).

The LLN for our model, given in [7], shows that under suitable assumptions on the parameters, the sequence of measure-valued processes  $\bar{A}^K = A^K/K$  converges as  $K \rightarrow \infty$  to a deterministic process  $\bar{A}^\infty$  in the Skorokhod space  $\mathbb{D}(\mathbb{R}_+, \mathcal{M}_+(\mathbb{R}_+))$ , where  $\mathcal{M}_+(\mathbb{R}_+)$  is the space of finite positive measures on  $\mathbb{R}_+$ . The limiting process is identified as the weak form of a generalised McKendrick-von Foerster Equation. In this paper, we establish the CLT (see Theorem 8) for the age structure, that is, the convergence of  $Z^K = \sqrt{K}(\bar{A}^K - \bar{A}^\infty)$  in an appropriate space, and identify the limit.

As usual, to show convergence we show tightness and uniqueness of the limit. The tightness is shown by using the Sobolev embedding approach and Aldous-Rebolledo tightness criteria, the method used in Bansaye et. al. [3], Meleard [19], and Tran [25]. Since  $Z^K$  is a signed measure-valued process, and the space of signed measures with the topology of weak convergence is not metrizable ([3],[19],[27]), we embed the space of signed measures in suitable Sobolev spaces (which are also Hilbert spaces), and apply Sobolev embedding techniques with some Hilbertian properties.

While the Sobolev embedding technique has been much used (e.g. [3], [4], [19]) since introduced by Metivier [21], and there are seminal papers in the field such as [3] and [16], our approach has a number of differences. We set up evolution equations for a branching process, fusing branching and stochastic analysis. This is done by using the Ulam-Harris representation. A simplifying technical feature of our model is that we can work on the bounded domain  $\mathbb{T}^* := [0, T + a^*]^1$ , where  $a^*$  is the age of the oldest individual alive at time 0, and this avoids the use of weighted Sobolev spaces (see page 4).

We show that the fluctuation process  $Z^K$  takes values in the space of finite signed measures, and give its evolution equation in an appropriate space ( $W^{-2}(\mathbb{T}^*)$ , see Subsection 3.3). In the limit (CLT), Fréchet derivatives of the rate functions naturally appear. They replace ordinary derivatives in the density-dependent case where dependence is on the total mass of the measure.

In Section 3 we state the results. The proof of the CLT is given in Section 4. In Section 6, we give some examples.

Throughout this paper, we use  $c$  with and without subscript to denote constants that may be different from line to line, but all independent of  $K$ .  $\mathbb{N}$  stands for the set of natural numbers and  $\mathbb{N}_0$  for the set of non-negative integers. For a Borel measure  $\mu$  on  $E$  and a measurable function  $f$ ,  $(f, \mu) := \int_E f(x) \mu(dx)$  and  $|\mu| := (1, \mu)$ . The Skorokhod space  $\mathbb{D}(\mathbb{T}, \mathcal{M})$  consists of all càdlàg functions from  $\mathbb{T} = [0, T]$  to  $\mathcal{M}$ . We will take  $\mathcal{M}$  to be a space of measures (for LLN) and the dual of a suitable Sobolev space of functions (for CLT).

---

<sup>1</sup> $T + a^*$  is an upper bound to the age of the oldest individual alive at time  $T$ .

## 2. EVOLUTION EQUATION AND SEMIMARTINGALE DECOMPOSITION

In this section we derive a semimartingale decomposition of the branching model. We shall adopt the classical, well-known in branching (e.g. [8]), Ulam-Harris labelling, as presented in [10] and developed in [11].

We use the set

$$I = \bigcup_{n=1}^{\infty} \mathbb{N}^n$$

to denote all possible individuals;  $\mathbb{N}$  corresponds to the possible individuals of the starting generation,  $\mathbb{N}^2$  corresponds to the possible individuals of the second generation, and so forth. We allow an arbitrary finite number of individuals at the start of the process at time  $t = 0$ . The individuals in the first (starting) generation are labelled  $1, 2, 3, \dots$ . For each individual  $y \in I$ , the children of  $y$  are consecutively labelled  $y1, y2, y3, \dots$  as they are born. Here  $yi$  is the concatenated vector of the coordinates of  $y \in I$  and  $i \in \mathbb{N}$ .

We assume that the age of each individual increases at rate 1 until the individual dies. Upon death it may split into a random number of offspring. During its lifetime the individual may give birth to a random number of offspring. The offspring generated in both situations are referred to as the children of the individual, and both situations are considered as births.

We denote by  $\tau_y$ ,  $\lambda_y$  and  $\sigma_y = \tau_y + \lambda_y$  respectively the time of birth, the life span and the death time of individual  $y$ . In particular, the maternal age for the birth of the  $j$ th child (during lifetime or by splitting at death) of individual  $y$  is  $\tau_{yj} - \tau_y$ . Also, if  $y$  has precisely  $n$  children, then

$$\tau_y < \tau_{y1} \leq \dots \leq \tau_{yn} \leq \sigma_y \text{ and } \tau_{y(n+1)} = \infty.$$

The population starts from an initial age distribution  $A_0$  with mass one at given ages  $x_1, x_2, \dots, x_{|A_0|}$ . Put  $\tau_i = -x_i$ ,  $i = 1, 2, \dots, |A_0|$  for the birth times of these ancestors (first generation).

The age distribution  $A_t$  at time  $t \geq 0$  allots a unit weight to the age  $(t - \tau_y)$  of each individual ( $y \in I$ ) that is alive at time  $t$ ,

$$A_t = \sum_{y \in I} \mathbf{1}_{\tau_y \leq t < \sigma_y} \delta_{t - \tau_y}. \quad (1)$$

For each  $t$ ,  $A_t$  is a finite discrete measure on  $\mathbb{R}_+$  and the collection  $(A_t)_{t \geq 0}$  is known as the age structure of the population.

Two processes that determine the evolution of population are the way the individuals enter and the way they exit. Denote by  $B(t)$  the number of individuals born by time  $t$ , and by  $D(x, t)$  the number of individuals who died by time  $t$  and whose life span was  $x$  or less, then

$$B(t) = \sum_{y \in I} \mathbf{1}_{\tau_y \leq t}, \quad D(x, t) = \sum_{y \in I} \mathbf{1}_{\lambda_y \leq x, \sigma_y \leq t}.$$

Before we give the fundamental equation for the evolution of the population, we make an important observation (which allows us to work on a bounded time interval and to avoid using weighted Sobolev spaces).

Recall that  $a^*$  is the age of the oldest individual in the starting generation, that is,

$$a^* = \inf\{x > 0 : A_0((x, \infty)) = 0\}.$$

Since we look at convergence on a finite time interval  $\mathbb{T}$ , the age of any individual at time  $t \leq T$  will not be more than  $T + a^*$ , thus the support of  $A_t$  is contained in  $[0, T + a^*]$ . Henceforth denote by  $T^* = T + a^*$ .

While our focus is indeed on functions of a single variable, the proof of the CLT requires a semimartingale decomposition for functions of two variables. Consequently, we consider test functions of two variables whose domain is limited to the bounded rectangle  $\mathbb{T}^* \times \mathbb{T}$ .

**Theorem 1.** *For any  $f \in C^{1,1}(\mathbb{T}^* \times \mathbb{T})$  and  $t \in \mathbb{T}$ , the age structure process  $A$  satisfies*

$$\begin{aligned} (f(\cdot, t), A_t) &= (f(\cdot, 0), A_0) + \int_0^t (\partial_1 f(\cdot, s) + \partial_2 f(\cdot, s), A_s) ds \\ &\quad + \int_{[0,t]} f(0, s) B(ds) - \int_{\mathbb{T}^* \times [0,t]} f(x, s) D(dx, ds). \end{aligned} \quad (2)$$

*Proof.* Note that  $(f(\cdot, t), A_t) = \sum_{y \in I} f(t - \tau_y, t) \mathbf{1}_{\tau_y \leq t < \sigma_y}$ . Let  $g(t) = f(t - \tau_y, t)$ , then  $g'(t) = \partial_1 f(t - \tau_y, t) + \partial_2 f(t - \tau_y, t)$  and

$$g(t) \mathbf{1}_{\tau_y \leq t < \sigma_y} - g(0) \mathbf{1}_{\tau_y < 0} = \int_0^t g'(s) \mathbf{1}_{\tau_y \leq s < \sigma_y} ds + g(\tau_y) \mathbf{1}_{0 \leq \tau_y \leq t} - g(\sigma_y) \mathbf{1}_{\sigma_y \leq t}.$$

Summing over  $y$ , we get (2).  $\square$

To arrive at compensators for the two processes in the RHS of (2), we assume the existence of birth and death rates, dependent on the age and also upon the population age structure (cf. [12]). The number of births by time  $t$  consists of births by living mothers and births by splitting,  $B = \check{B} + \widehat{B}$ . An individual aged  $x$  at time  $t$  gives birth at rate  $b_{A_t}(x)$  and dies at rate  $h_{A_t}(x)$ , allowing for multiple births.

Denote by  $\check{m}_{A_t}(x)$  and  $\widehat{m}_{A_t}(x)$  the conditional expected number of children at a bearing of a living mother aged  $x$  at time  $t$  and at splitting (i.e. death), respectively. Thus the conditional intensity of births of an individual aged  $x$  at time  $t$ , given the sigma-algebra  $\mathcal{F}_t$  of all events up to time  $t$  is  $\check{m}_{A_t}(x) b_{A_t}(x) + \widehat{m}_{A_t}(x) h_{A_t}(x)$ . We also denote the conditional second moment of the number of children at a bearing of a living mother aged  $x$  at time  $t$  by  $\check{v}_{A_t}(x)$ , and similarly the conditional second moment of the number of children at splitting by  $\widehat{v}_{A_t}(x)$ .

Let

$$\begin{aligned} Q^f(t) &= \int_{\mathbb{T}^* \times [0,t]} f(x, s) D(dx, ds), \\ \check{B}_t^f &= \int_{[0,t]} f(0, s) \check{B}(ds) \quad \text{and} \quad \widehat{B}_t^f = \int_{[0,t]} f(0, s) \widehat{B}(ds). \end{aligned}$$

Their compensators are given by the following result.

**Proposition 2.** *For every  $f \in C(\mathbb{T}^* \times \mathbb{T})$  and  $t \in \mathbb{T}$ ,*

$$\begin{aligned} H_t^f &:= \int_0^t (f(\cdot, s) h_{A_s}, A_s) ds, \\ \int_0^t f(0, s) (b_{A_s} \check{m}_{A_s}, A_s) ds, \quad \text{and} \quad \int_0^t f(0, s) (h_{A_s} \widehat{m}_{A_s}, A_s) ds \end{aligned}$$

*are the compensators of  $Q_t^f$ ,  $\check{B}_t^f$ , and  $\widehat{B}_t^f$  respectively.*

*Proof.* By the very definition of death rate, and the convention that all rates vanish for negative arguments,  $\mathbf{1}_{\sigma_y \leq t} - \int_0^{t \wedge \sigma_y} h_{A_s}(s - \tau_y) ds$  is a martingale. That is, for any bounded function  $g$ ,

$$\mathbb{E}[g(\sigma_y) \mathbf{1}_{\sigma_y > t} | \mathcal{F}_t] = \int_t^\infty g(s) \mathbb{E}[h_{A_s}(s - \tau_y) \mathbf{1}_{\sigma_y > s} | \mathcal{F}_t] ds$$

or equivalently that

$$\mathbb{P}(\sigma_y \in ds \cap (t, +\infty) | \mathcal{F}_t) = \mathbf{1}_{(t, +\infty)}(s) \mathbb{E}[h_{A_s}(s - \tau_y) \mathbf{1}_{\sigma_y > s} | \mathcal{F}_t] ds.$$

In particular

$$\lim_{\delta \downarrow 0} \frac{1}{\delta} \mathbb{P}(t < \sigma_y \leq t + \delta | \mathcal{F}_t) = h_{A_t}(t - \tau_y) \mathbf{1}_{\sigma_y > t}.$$

Now,  $Q_t^f = \sum_{y \in I} \mathbf{1}_{\sigma_y \leq t} f(\lambda_y, \sigma_y)$  is adapted to the filtration  $\mathcal{F} = \{\mathcal{F}_t\}$  and for any  $u > 0$ ,

$$\begin{aligned} \mathbb{E}[Q_{t+u}^f | \mathcal{F}_t] &= Q_t^f + \sum_{y \in I} \mathbb{E}[f(\sigma_y - \tau_y, \sigma_y) \mathbf{1}_{t < \sigma_y \leq t+u} | \mathcal{F}_t] \\ &= Q_t^f + \sum_{y \in I} \int_t^{t+u} \mathbb{E}[f(s - \tau_y, s) h_{A_s}(s - \tau_y) \mathbf{1}_{\sigma_y > s} | \mathcal{F}_t] ds; \end{aligned}$$

and similarly,  $H^f$  is adapted to  $\mathcal{F}$ , continuous and

$$\mathbb{E}[H_{t+u}^f | \mathcal{F}_t] = H_t^f + \sum_{y \in I} \mathbb{E}\left[\int_t^{t+u} f(s - \tau_y, s) h_{A_s}(s - \tau_y) \mathbf{1}_{\tau_y \leq s < \sigma_y} ds \middle| \mathcal{F}_t\right].$$

Hence,  $H^f$  is the compensator of  $Q^f$ , viewing that  $h_{A_s}(s - \tau_y) = 0$  for  $s < \tau_y$ . The proof for other compensators follows from the fact that  $\int_0^t (b_{A_s} \check{m}_{A_s}, A_s) ds$  and  $\int_0^t (h_{A_s} \hat{m}_{A_s}, A_s) ds$  are compensators for  $\check{B}$  and  $\hat{B}$ .  $\square$

Having found the compensators we identify the relevant martingales. The proof of the following proposition is standard and is therefore omitted.

**Proposition 3.** *The following processes are martingales*

$$\begin{aligned} M_{D,f}(t) &:= \int_{\mathbb{T}^* \times [0,t]} f(x, s) D(dx, ds) - \int_0^t (f(\cdot, s) h_{A_s}, A_s) ds \\ M_{\check{B},f}(t) &:= \int_{[0,t]} f(0, s) \check{B}(ds) - \int_0^t f(0, s) (b_{A_s} \check{m}_{A_s}, A_s) ds \\ M_{\hat{B},f}(t) &:= \int_{[0,t]} f(0, s) \hat{B}(ds) - \int_0^t f(0, s) (h_{A_s} \hat{m}_{A_s}, A_s) ds \end{aligned}$$

with predictable quadratic variations

$$\begin{aligned} \langle M_{D,f} \rangle_t &= \int_0^t (f^2(\cdot, s) h_{A_s}, A_s) ds, \quad \langle M_{\check{B},f} \rangle_t = \int_0^t f^2(0, s) (b_{A_s} \check{v}_{A_s}, A_s) ds, \\ \langle M_{\hat{B},f} \rangle_t &= \int_0^t f^2(0, s) (h_{A_s} \hat{v}_{A_s}, A_s) ds. \end{aligned}$$

We combine the rates  $n = b\check{m} + h\hat{m}$  and  $w = b\check{v} + h\hat{v}$ , and also the martingales. From the basic equation (2) we obtain the following semimartingale decomposition.

**Corollary 4.** For  $t \in \mathbb{T}$  and  $f \in C^{1,1}(\mathbb{T}^* \times \mathbb{T})$ ,

$$(f(\cdot, t), A_t) = (f(\cdot, 0), A_0) + \int_0^t (\partial_1 f(\cdot, s) + \partial_2 f(\cdot, s) - f(\cdot, s)h_{A_s} + f(0, s)n_{A_s}, A_s)ds + M_t^f, \quad (3)$$

where  $M_t^f$  is a locally-square-integrable  $\{\mathcal{F}_t\}$ -martingale with predictable quadratic variation

$$\langle M^f \rangle_t = \int_0^t (f^2(0, s)w_{A_s} + h_{A_s}f^2(\cdot, s) - 2f(0, s)h_{A_s}\widehat{m}_{A_s}f(\cdot, s), A_s)ds. \quad (4)$$

*Proof.* Here, we prove (4). Note that the martingales  $M_{D,f}$ ,  $M_{\check{B},f}$  and  $M_{\widehat{B},f}$  are purely discontinuous. Since  $M_{\check{B},f}$  and  $M_{\widehat{B},f}$  do not jump together,  $[M_{\check{B},f}, M_{\widehat{B},f}]_t$  and thus  $\langle M_{\check{B},f}, M_{\widehat{B},f} \rangle_t$  are zero. Similarly for  $M_{D,f}$  and  $M_{\check{B},f}$ , giving  $\langle M_{D,f}, M_{\check{B},f} \rangle_t = 0$ . However,  $M_{D,f}$  and  $M_{\widehat{B},f}$  jump together when there is a birth by splitting with  $\Delta M_{D,f}(t) = \sum_{y \in I} f(\lambda_y, t)\mathbf{1}_{\sigma_y=t}$  and  $\Delta M_{\widehat{B},f}(t) = \sum_{y \in I} f(0, t)\mathbf{1}_{\sigma_y=t} \sum_{i \in \mathbb{N}} \mathbf{1}_{\tau_{yi}=t}$ . Thus,

$$[M_{D,f}, M_{\widehat{B},f}]_t = \sum_{s \leq t} \sum_{y \in I} f(0, s)f(\lambda_y, s)\mathbf{1}_{\sigma_y=s} \sum_{i \in \mathbb{N}} \mathbf{1}_{\tau_{yi}=s}$$

and its compensator

$$\langle M_{D,f}, M_{\widehat{B},f} \rangle_t = \int_0^t f(0, s)(f(\cdot, s)h_{A_s}\widehat{m}_{A_s}, A_s)ds.$$

Finally, as  $M_t^f = M_{\check{B},f}(t) + M_{\widehat{B},f}(t) - M_{D,f}(t)$ , we have

$$\langle M^f \rangle_t = \langle M_{\check{B},f} \rangle_t + \langle M_{\widehat{B},f} \rangle_t + \langle M_{D,f} \rangle_t - 2\langle M_{D,f}, M_{\widehat{B},f} \rangle_t,$$

which gives (4).  $\square$

In particular, taking  $f$  as a function of the first variable  $x$  only, we recover Equation (2.6) of [12].

**Corollary 5.** For  $t \in \mathbb{T}$  and  $f \in C^1(\mathbb{T}^*)$ ,

$$(f, A_t) = (f, A_0) + \int_0^t (L_{A_s}f, A_s)ds + M_t^f, \quad (5)$$

where

$$L_A f = f' - h_A f + f(0)n_A$$

and  $M_t^f$  is a locally-square-integrable martingale with predictable quadratic variation

$$\langle M^f \rangle_t = \int_0^t (f^2(0)w_{A_s} + h_{A_s}f^2 - 2f(0)h_{A_s}\widehat{m}_{A_s}f, A_s)ds. \quad (6)$$

**Remark 6.** We can also obtain the predictable quadratic covariation of the martingale with two test functions. For  $f, g \in C(\mathbb{T}^* \times \mathbb{T})$  and  $t \in \mathbb{T}$ ,

$$\begin{aligned} \langle M^f, M^g \rangle_t = \int_0^t & \left( f(0, s)g(0, s)w_{A_s} + h_{A_s}f(\cdot, s)g(\cdot, s) \right. \\ & \left. - h_{A_s}\widehat{m}_{A_s}(f(0, s)g(\cdot, s) + g(0, s)f(\cdot, s)), A_s \right) ds. \end{aligned}$$

3. CLT FOR BRANCHING PROCESSES DEPENDENT ON PARAMETER  $K$ 

We now look at the case of a branching process dependent on some (large) index  $K$ .  $K$  may, for example, represent the population carrying capacity, a threshold below which the process is supercritical and above which it is subcritical. The notion of carrying capacity plays a great role in biological population dynamics.

We are interested in the approximation of such a process for large  $K$  and are therefore lead to consider a family of branching processes indexed by  $K$ . All objects introduced in the previous sections will now carry the extra label  $K$ :  $A^K$ ,  $b^K$ ,  $h^K$  etc.

Throughout the remainder of the paper, we make one simplifying (and reasonable) assumption in that the ages of all individuals in all starting generations are bounded. We denote (with a slight abuse of notation) by  $a^*$  “the age of the oldest individual” at  $t = 0$

$$a^* := \sup_{K \geq 1} \left( \inf\{x > 0 : A_0^K((x, \infty)) = 0\} \right) < \infty.$$

As before,  $\mathbb{T}^* = [0, T^*]$  with  $T^* = T + a^*$ . For each  $K$ ,  $A^K = (A_t^K)_{t \in \mathbb{T}}$  is a càdlàg positive measure-valued process on  $\mathbb{T}^*$ , i.e.  $(A_t^K)_{t \in \mathbb{T}} \in \mathbb{D}(\mathbb{T}, \mathcal{M}_+(\mathbb{T}^*))$ . Without loss of generality, we assume that  $A_0^K$  is deterministic.

As we shall focus on situations where  $\bar{A}^K := A^K/K$  converges to a non-degenerate limit, a new parametrisation of the rates is needed, one that involves  $\bar{A}^K$  rather than  $A^K$  itself.

We have, immediately from Equation (5), the following evolution of  $\bar{A}^K$ :

$$(f, \bar{A}_t^K) = (f, \bar{A}_0^K) + \int_0^t (L_{\bar{A}_s^K}^K f, \bar{A}_s^K) ds + \frac{1}{K} M_t^{f,K}, \quad (7)$$

where

$$L_A^K f = f' - h_A^K f + f(0)n_A^K \quad (8)$$

and  $M_t^{f,K}$  is a martingale. A similar representation with functions of two variables is also used later in proofs.

**3.1. The Law of Large Numbers.** The LLN was established in [7] under the following conditions, referred to as *smooth demography*:

- (C0) The model parameters  $b$ ,  $h$ ,  $\hat{m}$ ,  $\check{m}$ ,  $\hat{v}$  and  $\check{v}^2$  are uniformly bounded, that is,  $\sup_{K,A,x} b_A^K(x) < \infty$ , et cetera. Note that the supremum with respect to  $A$  is taken over  $A \in \mathcal{M}_+(\mathbb{T}^*)$ .
- (C1) The model parameters  $b$ ,  $h$  and  $m$  are normed uniformly Lipschitz in the following sense: there is a  $c > 0$  such that for all  $K \geq 1$ ,  $\|b_A^K - b_B^K\|_\infty \leq c\|A - B\|$ , where  $\|\mu - \nu\| := \sup_{\|f\|_\infty \leq 1} |(f, \mu - \nu)|$ ; the same applies to  $h$  and  $m$ .
- (C2) The limit (pointwise in  $A$  and uniform in  $x$ )  $\lim b_A^K =: b_A^\infty$  exists; the same applies to limits  $\lim h_A^K =: h_A^\infty$  and  $\lim m_A^K =: m_A^\infty$ .
- (C3)  $\bar{A}_0^K \Rightarrow \bar{A}_0^\infty$ ,  $\sup_K |\bar{A}_0^K| < \infty$ .

We remark that in [7], the Prokhorov metric is used for (C1). However, since we shall work in spaces  $C^{-j}$  and  $W^{-j}$  (see Section 3.3) for the CLT, it is more natural to use the norms in these spaces. The norm  $\|\cdot\|$  is precisely  $\|\cdot\|_{C^{-0}}$  defined in Section 3.3. It can be shown that the LLN remains valid with this (C1).

<sup>2</sup>The qualifiers  $\hat{\cdot}$  and  $\check{\cdot}$  will be dropped in any statement that refers to either parameter.

**Theorem 7** ([7]). *Under the smooth demography condition, as  $K \rightarrow \infty$ ,  $\bar{A}^K$  converges weakly in the Skorokhod space  $\mathbb{D}(\mathbb{T}, \mathcal{M}_+(\mathbb{T}^*))$  to the limiting process  $\bar{A}^\infty$ , which is deterministic and satisfies, for  $f \in C^1(\mathbb{T}^*)$  and  $t \in \mathbb{T}$ ,*

$$\begin{aligned} (f, \bar{A}_t^\infty) &= (f, \bar{A}_0^\infty) + \int_0^t (f' - h_{\bar{A}_s^\infty}^\infty f + f(0)n_{\bar{A}_s^\infty}^\infty, \bar{A}_s^\infty) ds \\ &= (f, \bar{A}_0^\infty) + \int_0^t (L_{\bar{A}_s^\infty}^\infty f, \bar{A}_s^\infty) ds, \end{aligned} \quad (9)$$

where  $n_{\bar{A}_s^\infty}^\infty = b_{\bar{A}_s^\infty}^\infty \check{m}_{\bar{A}_s^\infty}^\infty + h_{\bar{A}_s^\infty}^\infty \hat{m}_{\bar{A}_s^\infty}^\infty$ , and  $L_{\bar{A}_s^\infty}^\infty f = f' - h_{\bar{A}_s^\infty}^\infty f + f(0)n_{\bar{A}_s^\infty}^\infty$ .

It follows by the Monotone Class Theorem (e.g. [5, I.22.1]) that (9) also holds for test functions of two variables,  $f \in C^{1,1}(\mathbb{T}^* \times \mathbb{T})$ , and  $t \in \mathbb{T}$ . This fact will be used later in the representation of the fluctuation process.

As remarked in [7], if  $\bar{A}_0^\infty$  has a density, then  $\bar{A}_t^\infty$  has a density. Call it  $a(x, t)$ . In this case, Equation (9) is the weak form of the McKendrick-von Foerster equation for the density:

$$\left( \frac{\partial}{\partial x} + \frac{\partial}{\partial t} \right) a(x, t) = -a(x, t) h_{\bar{A}_t^\infty}^\infty(x), \quad a(0, t) = \int_0^{t+a^*} n_{\bar{A}_t^\infty}^\infty(x) a(x, t) dx.$$

**3.2. The fluctuation process  $Z^K$ .** For each  $t$  and  $K$ ,  $Z_t^K := \sqrt{K}(\bar{A}_t^K - \bar{A}_t^\infty)$  is a finite signed measure that, in view of (7) and (9), can be represented as

$$\begin{aligned} (f, Z_t^K) &= (f, Z_0^K) + \sqrt{K} \int_0^t \left( L_{\bar{A}_s^K}^K f - L_{\bar{A}_s^\infty}^\infty f, \bar{A}_s^\infty \right) ds \\ &\quad + \int_0^t \left( L_{\bar{A}_s^K}^K f, Z_s^K \right) ds + \tilde{M}_t^{f,K}, \end{aligned} \quad (10)$$

where  $\tilde{M}_t^{f,K}$  is a martingale with predictable quadratic variation

$$\langle \tilde{M}^{f,K} \rangle_t = \int_0^t (f^2(0)w_{\bar{A}_s^K}^K + h_{\bar{A}_s^K}^K f^2 - 2f(0)h_{\bar{A}_s^K}^K \hat{m}_{\bar{A}_s^K}^K f, \bar{A}_s^K) ds.$$

**3.3. Relevant spaces and embeddings.** Let  $C^j(\mathbb{T}^*)$ ,  $j \in \mathbb{N}_0$ , denote the space of continuous functions on  $\mathbb{T}^*$  with continuous derivatives up to order  $j$ . Since  $\mathbb{T}^*$  is a bounded domain, the functions in  $C^j(\mathbb{T}^*)$  as well as their  $j$  derivatives are bounded with the norm

$$\|f\|_{C^j(\mathbb{T}^*)} = \max_{0 \leq i \leq j} \sup_{x \in \mathbb{T}^*} |f^{(i)}(x)|.$$

The Sobolev space  $W^j(\mathbb{T}^*)$  is the closure of  $C^\infty(\mathbb{T}^*)$  with respect to the norm

$$\|f\|_{W^j(\mathbb{T}^*)} = \left( \sum_{i=0}^j \int_{\mathbb{T}^*} (f^{(i)}(x))^2 dx \right)^{1/2},$$

where  $f^{(i)}$  is the (weak) derivative of  $f$  (see e.g. [1]). The space  $W^j(\mathbb{T}^*)$  is a Hilbert space with inner product  $\langle f, g \rangle_{W^j(\mathbb{T}^*)} = \sum_{i=0}^j \int_{\mathbb{T}^*} f^{(i)}(x) g^{(i)}(x) dx$ .

For the rest of this paper, we assume, unless otherwise specified, that function are defined on the domain  $\mathbb{T}^*$  and suppress the label  $\mathbb{T}^*$ ; e.g.  $W^j$  means  $W^j(\mathbb{T}^*)$ .

The following embeddings hold

$$C^j \hookrightarrow W^j, \quad W^{j+1} \hookrightarrow C^j \quad \text{and} \quad W^{j+1} \xhookrightarrow{H.S.} W^j,$$



where H.S. stands for Hilbert-Schmidt embedding. Let  $C^{-j}(\mathbb{T}^*)$  and  $W^{-j}(\mathbb{T}^*)$  denote the dual spaces of, respectively  $C^j(\mathbb{T}^*)$  and  $W^j(\mathbb{T}^*)$ . Then,

$$W^{-j} \hookrightarrow C^{-j}, \quad C^{-j} \hookrightarrow W^{-(j+1)} \quad \text{and} \quad W^{-j} \xhookrightarrow{\text{H.S.}} W^{-(j+1)}.$$

In particular, we have

$$C^{-0} \hookrightarrow W^{-2} \xhookrightarrow{\text{H.S.}} W^{-3} \xhookrightarrow{\text{H.S.}} W^{-4}.$$

A consequence of the Hilbert-Schmidt embedding is that the unit ball in  $W^{-3}$  is compact in  $W^{-4}$ , a fact exploited in the proof of the componentwise tightness of the fluctuation processes (see proof of Theorem 11).

For each  $t$  and  $K$ ,  $Z_t^K$ , as a signed measure, belongs to  $C^{-0}$ , and when considered as an element of  $C^{-1}$  (or any  $W^{-j}$ ,  $j \geq 2$ ) admits representation (10). The embedding  $W^{-3} \hookrightarrow W^{-4}$  is used to show tightness of the family  $Z^K$ , seen as processes in  $W^{-4}$ , whereas the embedding  $W^{-2} \hookrightarrow W^{-3}$  is used to bound  $\mathbb{E}[\sup_{t \leq T} \|Z_t^K\|_{W^{-3}}]$ , which in turn is also used to prove tightness.

We shall use the following general results, the proofs of which are standard and are therefore omitted. For any  $f \in C^j$  and  $g \in W^j$ ,  $j \in \mathbb{N}_0$ ,

$$\|fg\|_{W^j} \leq c \|f\|_{C^j} \|g\|_{W^j}. \quad (11)$$

Let  $(p_l^j)_{l \geq 1}$  denote a complete orthonormal basis of  $W^j$ ,  $j \in \mathbb{N}$ . Then, for any  $x_1 \in \mathbb{T}^*$  and  $x_2 \in \mathbb{T}^*$ ,

$$\left| \sum_{l \geq 1} p_l^j(x_1) p_l^j(x_2) \right| \leq c. \quad (12)$$

**3.4. Statement of the CLT.** Further to (C0)-(C3), we shall make the following assumptions.

- (A0) Conditions (C1) and (C2) hold also for  $v$ .
- (A1) The random variables  $\check{\xi}_A^K(x)$  and  $\hat{\xi}_A^K(x)$  (the number of offspring by a living mother and by splitting at death) are dominated by a random variable  $\Xi$  with finite mean and variance.
- (A2) The reproduction parameters  $b_A^K(x)$ ,  $h_A^K(x)$  and  $m_A^K(x)$  and their limits (in the sense of (C2)) are in  $C^4$ , in the argument  $x$ , with convergence in  $C^4$ . Moreover,  $\sqrt{K} \sup_A \|b_A^K - b_A^\infty\|_\infty \rightarrow 0$  as  $K \rightarrow \infty$ ,  $\sup_{K,A} \|b_A^K\|_{C^3} < \infty$  and  $\sup_A \|b_A^\infty\|_{C^4} < \infty$ ; similarly for parameters  $h$  and  $m$ .
- (A3) The limiting parameters (as functions of  $A$ ) are Fréchet differentiable at every  $A$ . Namely, for every  $A_0$ , there exists a continuous linear operator  $\partial_A b_{A_0}^\infty : W^{-4} \rightarrow L_\infty$  such that

$$\lim_{\|B\|_{W^{-4}} \rightarrow 0} \frac{1}{\|B\|_{W^{-4}}} \|b_{A_0+B}^\infty - b_{A_0}^\infty - \partial_A b_{A_0}^\infty(B)\|_\infty = 0.$$

Moreover,  $\sup_{A_0} \|\partial_A b_{A_0}^\infty\|_{\mathbb{L}^{-4}} \leq c$ , where  $\mathbb{L}^{-4} = L(W^{-4}, L_\infty)$  denotes the space of continuous linear mappings from  $W^{-4}$  to  $L_\infty$ . The same applies to parameters  $h^\infty$  and  $m^\infty$ .

- (A4)  $Z_0^K$  converges to  $Z_0^\infty$  in  $W^{-4}$  and  $\sup_K \|Z_0^K\|_{W^{-2}} < \infty$ .

**Theorem 8.** *Assume (A0)-(A4) in addition to the smooth demography condition (C0)-(C3). Then, as  $K \rightarrow \infty$ , the process  $(Z_t^K)_{t \in \mathbb{T}}$  converges weakly in  $\mathbb{D}(\mathbb{T}, W^{-4})$*

to  $(Z_t^\infty)_{t \in \mathbb{T}}$  satisfying the equation, for  $f \in W^4$ ,

$$\begin{aligned} (f, Z_t) &= (f, Z_0) + \int_0^t (-\partial_A h_{\bar{A}_s}^\infty(Z_s)f + f(0)\partial_A n_{\bar{A}_s}^\infty(Z_s), \bar{A}_s^\infty) ds \\ &\quad + \int_0^t (f' - h_{\bar{A}_s}^\infty f + f(0)n_{\bar{A}_s}^\infty, Z_s) ds + \tilde{M}_t^{f, \infty} \end{aligned} \quad (13)$$

where  $n_{\bar{A}_s}^\infty = b_{\bar{A}_s}^\infty \check{m}_{\bar{A}_s}^\infty + h_{\bar{A}_s}^\infty \hat{m}_{\bar{A}_s}^\infty$  and  $\tilde{M}_t^{f, \infty}$  is a continuous Gaussian martingale with predictable quadratic variation

$$\langle \tilde{M}^{f, \infty} \rangle_t = \int_0^t \left( f^2(0)w_{\bar{A}_s}^\infty + h_{\bar{A}_s}^\infty f^2 - 2f(0)h_{\bar{A}_s}^\infty \hat{m}_{\bar{A}_s}^\infty f, \bar{A}_s^\infty \right) ds,$$

with  $w_{\bar{A}_s}^\infty = b_{\bar{A}_s}^\infty \check{v}_{\bar{A}_s}^\infty + h_{\bar{A}_s}^\infty \hat{v}_{\bar{A}_s}^\infty$ .

**Corollary 9 (SPDE).** *The limiting process  $(Z_t^\infty)_{t \in \mathbb{T}}$  satisfies the following SPDE:*

$$\begin{aligned} dZ_t(dx) &= -\partial_A h_{\bar{A}_t}^\infty(Z_t) \bar{A}_t^\infty(dx) dt + (\partial_A n_{\bar{A}_t}^\infty(Z_t), \bar{A}_t^\infty) dt \delta_0(dx) \\ &\quad - (Z_t)'(dx) dt - h_{\bar{A}_t}^\infty Z_t(dx) dt + (n_{\bar{A}_t}^\infty, Z_t) dt \delta_0(dx) + d\tilde{M}_t^\infty(dx), \end{aligned}$$

where  $\tilde{M}^\infty$  is a Gaussian martingale measure such that  $(f, \tilde{M}_t^\infty) = \tilde{M}_t^{f, \infty}$ , and  $(Z_t)'$  is defined by  $(f, (Z_t)') = -(f', Z_t)$ .

**Proposition 10.** *Suppose that  $\partial_A h_{A_0}^\infty(B)(x)$  has the form  $(g^h(A_0, x, \cdot), B)$  for some  $g^h(A, x, \cdot) \in W^4$  with  $\sup_{A, x} \|g^h(A, x, \cdot)\|_{W^4} < \infty$ , and similarly  $\partial_A n_{A_0}^\infty(B)(x)$  is of the form  $(g^n(A_0, x, \cdot), B)$  for some  $g^n(A, x, \cdot) \in W^4$  with  $\sup_{A, x} \|g^n(A, x, \cdot)\|_{W^4} < \infty$ . Then,  $\nu_t : f \mapsto \mathbb{E}[(f, Z_t)]$  is a measure.*

We postpone the proofs of Corollary 9 and Proposition 10 to Section 5.

#### 4. PROOF OF CLT

We establish the tightness of the sequence  $Z^K$  first, and then show the uniqueness of the limit.

**4.1. Tightness of  $Z^K$ .** First we prove a result for the tightness of  $W^{-j}$ -valued processes in Skorokhod space  $\mathbb{D}(\mathbb{T}, W^{-j})$ , which we apply to  $Z_t^K$ .

**Theorem 11.** *Suppose  $(\mu^K)_{K \geq 1}$  is a sequence of  $W^{-j}$ -valued càdlàg processes. We assume that the dynamics of  $\mu^K$  are given by*

$$(f, \mu_t^K) = (f, \mu_0^K) + \int_0^t \Lambda_s^K f ds + \tilde{M}_t^{f, K}, \quad (14)$$

where  $\tilde{M}^{f, K}$  is a martingale with predictable quadratic variation of the form

$$\langle \tilde{M}^{f, K} \rangle_t = \int_0^t \Gamma_s^K f ds, \quad (15)$$

and  $\Lambda_t^K$  and  $\Gamma_t^K$  are functionals on  $W^j$ . The sequence  $(\mu^K)_{K \geq 1}$  is tight in  $\mathbb{D}(\mathbb{T}, W^{-j})$  if the following conditions are satisfied:

(T1) *There exists  $i < j$  such that for all  $t \in \mathbb{T}$ ,*

$$\sup_{K \geq 1} \mathbb{E}[\|\mu_t^K\|_{W^{-i}}] < \infty.$$

(T2) *There exists  $K_0 \geq 1$  such that*

$$\sup_{K \geq K_0} \mathbb{E} \left[ \sup_{t \leq T} \|\Lambda_t^K\|_{W^{-j}} \right] \leq c_T \quad (\text{i})$$

$$\sup_{K \geq K_0} \mathbb{E} \left[ \sup_{t \leq T} \left| \sum_{l \geq 1} \Gamma_t^K p_l^j \right| \right] \leq c_T. \quad (\text{ii})$$

We prove it by showing that the Aldous-Rebolledo criteria for tightness, stated below, holds. For more details see for example [14, pp. 34-35], [2].

**Lemma 12** (Aldous-Rebolledo). *Let  $H$  be a separable Hilbert space. A sequence  $(\mu^K)_{K \geq 1}$  of  $H$ -valued càdlàg processes is tight in  $\mathbb{D}(\mathbb{T}, H)$  if the following conditions are satisfied:*

- (A) *For every  $t \in \mathbb{T}$ ,  $(\mu_t^K)_{K \geq 1}$  is tight in  $H$ .*
- (B) *For each  $\epsilon_1, \epsilon_2 > 0$ , there exist  $\delta > 0$  and  $K_0 \geq 1$  such that for every sequence of stopping times  $\tau^K \leq T$ ,*

$$\sup_{K > K_0} \sup_{\zeta < \delta} \mathbb{P}(\|\mu_{(\tau^K + \zeta) \wedge T}^K - \mu_{\tau^K}^K\|_H > \epsilon_1) < \epsilon_2.$$

*If  $\mu_t^K$  admits a semimartingale decomposition, then for (B), it is sufficient to have it for the finite variation part and the predictable quadratic variation of the martingale part.*

*Proof of Theorem 11.* Note that, for  $i < j$ ,  $W^{-i} \xrightarrow[H.S.]{} W^{-j}$ , thus, the closed ball  $B_{W^{-i}}(R) := \{\mu \in W^{-i} : \|\mu\|_{W^{-i}} \leq R\}$  is compact in  $W^{-j}$ . Also,

$$\mathbb{P}(\mu_t^K \notin B_{W^{-i}}(R)) = \mathbb{P}(\|\mu_t^K\|_{W^{-i}} > R) \leq \frac{1}{R} \mathbb{E}[\|\mu_t^K\|_{W^{-i}}].$$

Therefore, if (T1) holds, there exists a compact set  $\mathcal{C}_\epsilon$  such that  $\mathbb{P}(\mu_t^K \notin \mathcal{C}_\epsilon) < \epsilon$  for all  $K$ , which in turn implies (A).

Next, we show that (T2) implies (B). Since  $\mu^K$  has the form  $\mu_t^K = V_t^K + \tilde{M}_t^K$ , it remains to show (B) for  $V_t^K$  and predictable quadratic variation  $\langle \langle M^K \rangle \rangle_t^3$ .

To see (B) for  $V_t^K$ , observe that by (14)

$$\begin{aligned} |(f, V_{(\tau^K + \zeta) \wedge T}^K - V_{\tau^K}^K)| &= \left| \int_{\tau^K}^{(\tau^K + \zeta) \wedge T} \Lambda_s^K f ds \right| \\ &\leq \int_{\tau^K}^{(\tau^K + \zeta) \wedge T} |\Lambda_s^K f| ds \leq \int_{\tau^K}^{(\tau^K + \zeta) \wedge T} \|\Lambda_s^K\|_{W^{-j}} \|f\|_{W^j} ds. \end{aligned}$$

Hence

$$\begin{aligned} \|V_{(\tau^K + \zeta) \wedge T}^K - V_{\tau^K}^K\|_{W^{-j}} &\leq \int_{\tau^K}^{(\tau^K + \zeta) \wedge T} \|\Lambda_s^K\|_{W^{-j}} ds \\ &= \int_0^\zeta \|\Lambda_{(\tau^K + s) \wedge T}^K\|_{W^{-j}} ds \leq \delta \sup_{t \leq T} \|\Lambda_t^K\|_{W^{-j}}. \end{aligned}$$

(B) now follows from condition (T2)(i) by Markov's inequality.

Write  $p_l$  for  $p_l^j$ . Since  $\|\tilde{M}_t^K\|_{W^{-j}}^2 = \sum_{l \geq 1} (\tilde{M}_t^{p_l, K})^2$  by the Riesz Representation Theorem and Parseval's identity, and for each  $l$ ,  $((\tilde{M}_t^{p_l, K})^2 - \langle \tilde{M}^{p_l, K} \rangle_t)_{t \in \mathbb{T}}$  is a

---

<sup>3</sup> $\langle \langle \tilde{M}^K \rangle \rangle_t$  is defined such that  $(\|\tilde{M}_t^K\|_{W^{-j}}^2 - \langle \langle \tilde{M}^K \rangle \rangle_t)_{t \in \mathbb{T}}$  is a martingale.

martingale, thus we have  $\langle \langle \tilde{M}^K \rangle \rangle_t = \sum_{l \geq 1} \langle \tilde{M}^{p_l, K} \rangle_t$ . To see (B) for  $\langle \langle \tilde{M}^K \rangle \rangle_t$ , by (15),

$$\begin{aligned} \left| \langle \langle \tilde{M}^K \rangle \rangle_{(\tau^K + \zeta) \wedge T} - \langle \langle \tilde{M}^K \rangle \rangle_{\tau^K} \right| &= \left| \sum_{l \geq 1} \langle \tilde{M}^{p_l, K} \rangle_{(\tau^K + \zeta) \wedge T} - \sum_{l \geq 1} \langle \tilde{M}^{p_l, K} \rangle_{\tau^K} \right| \\ &= \left| \sum_{l \geq 1} \int_{\tau^K}^{(\tau^K + \zeta) \wedge T} \Gamma_s^K p_l ds \right| = \left| \sum_{l \geq 1} \int_0^\zeta \Gamma_{(\tau^K + s) \wedge T}^K p_l ds \right| \end{aligned}$$

and by taking expectation,

$$\begin{aligned} \mathbb{E} \left[ \left| \langle \langle \tilde{M}^K \rangle \rangle_{(\tau^K + \zeta) \wedge T} - \langle \langle \tilde{M}^K \rangle \rangle_{\tau^K} \right| \right] &\leq \int_0^\zeta \mathbb{E} \left[ \left| \sum_{l \geq 1} \Gamma_{(\tau^K + s) \wedge T}^K p_l \right| \right] ds \\ &\leq \int_0^\zeta \mathbb{E} \left[ \sup_{t \leq T} \left| \sum_{l \geq 1} \Gamma_t^K p_l \right| \right] ds \leq \delta \mathbb{E} \left[ \sup_{t \leq T} \left| \sum_{l \geq 1} \Gamma_t^K p_l \right| \right]. \end{aligned}$$

(B) now follows from condition (T2)(ii) by Markov's inequality.  $\square$

The rest of the proof consists of checking conditions (T1) and (T2) in space  $W^{-4}$ . The proof is involved and requires somewhat different representations for  $Z_t^K$ , and is split into sections.

**4.2. Representation for  $Z_t^K$ .** As representation (10) involves the unbounded derivative operator ( $f \rightarrow f'$ ), we extend (10) to functions of two variables  $f(x, s)$  and apply the extension to the special case  $f(x, s) = \phi(x + t - s)$  (for some fixed  $t$  and some function  $\phi$ ). This results in the removal of the derivative operator but introduces a new challenge, that of representing the martingales as integrals with respect to a common measure.

From (7) and (9), we have, for test function of two variables  $f \in C^{1,1}(\mathbb{T}^* \times \mathbb{T})$  and  $t \in \mathbb{T}$ ,

$$\begin{aligned} (f(\cdot, t), Z_t^K) &= (f(\cdot, 0), Z_0^K) \\ &\quad + \sqrt{K} \int_0^t \left( -(h_{\bar{A}_s^K}^K - h_{\bar{A}_s^\infty}^\infty) f(\cdot, s) + f(0, s) (n_{\bar{A}_s^K}^K - n_{\bar{A}_s^\infty}^\infty), \bar{A}_s^\infty \right) ds \\ &\quad + \int_0^t \left( \partial_1 f(\cdot, s) + \partial_2 f(\cdot, s) - h_{\bar{A}_s^K}^K f(\cdot, s) + f(0, s) n_{\bar{A}_s^K}^K, Z_s^K \right) ds + \tilde{M}_t^{f, K}, \end{aligned} \quad (16)$$

where  $\tilde{M}_t^{f, K}$  is a martingale with predictable quadratic variation

$$\langle \tilde{M}^{f, K} \rangle_t = \int_0^t (f^2(0, s) w_{\bar{A}_s^K}^K + h_{\bar{A}_s^K}^K f^2(\cdot, s) - 2f(0, s) h_{\bar{A}_s^K}^K \hat{m}_{\bar{A}_s^K}^K f(\cdot, s), \bar{A}_s^K) ds. \quad (17)$$

As explained above, applying (16) to

$$f_s(x) \equiv f(x, s) = \phi(x + t - s) =: \Theta_{t-s} \phi(x)$$

(for a fixed  $t$ ) makes the term  $\partial_1 f(\cdot, s) + \partial_2 f(\cdot, s)$  vanish.

Next we obtain a representation for the corresponding martingale  $\tilde{M}_t^{f, K}$ .

Define the measure  $M_t$  as

$$\begin{aligned} M_t(dx) = & \delta_0(dx) \left( \check{B}([0, t]) - \int_0^t (b_{A_s} \check{m}_{A_s}, A_s) ds \right) \\ & + \delta_0(dx) \left( \hat{B}([0, t]) - \int_0^t (h_{A_s} \hat{m}_{A_s}, A_s) ds \right) \\ & - \left( \sum_{y \in I} \delta_{\lambda_y}(dx) \mathbf{1}_{\sigma_y \leq t} - \int_0^t A_s^h(dx) ds \right), \end{aligned} \quad (18)$$

where

$$A_t^h(dx) = \sum_{y \in I} \delta_{t-\tau_y}(dx) h_{A_t}(x) \mathbf{1}_{\tau_y \leq t < \sigma_y}.$$

Then by direct calculations we can see that the martingale  $M_t^f$  in (5) is precisely the integral of  $f$  with respect to  $M_t$ , i.e.  $M_t^f = (f, M_t)$ . It is easy to extend the definition of the integral to functions of two variables  $f \in C(\mathbb{T}^* \times \mathbb{T})$  so that  $\int_0^t (f(\cdot, s), dM_s)$  coincides with  $M_t^f$  in (3).

Indeed, since  $((g, M_t))_{t \in \mathbb{T}}$  is a martingale for any  $g \in C(\mathbb{T}^*)$ , thus, for any  $\varphi \in C(\mathbb{T})$ , the integral  $\int_0^t \varphi(s) d(g, M_s)$ ,  $t \in \mathbb{T}$ , is a well-defined martingale with predictable quadratic variation

$$\left\langle \int_0^\cdot \varphi(s) d(g, M_s) \right\rangle_t = \int_0^t \varphi^2(s) (g^2(0) w_{A_s} + h_{A_s} g^2 - 2g(0) h_{A_s} \hat{m}_{A_s} g, A_s) ds.$$

Write  $\int_0^t (\varphi(s) g, dM_s)$  for  $\int_0^t \varphi(s) d(g, M_s)$ . The extension to an arbitrary  $f \in C(\mathbb{T}^* \times \mathbb{T})$  is obtained by the usual application of the Monotone Class Theorem (e.g. [5, I.22.1]).

Let  $\tilde{M}^K = \frac{1}{\sqrt{K}} M^K$ . Since, for a fixed  $t \in \mathbb{T}$ , the function  $f(x, s) = \Theta_{t-s} \phi(x)$  satisfies  $f(x, t) = \phi(x)$ , (16) reduces to (19) below.

**Corollary 13.** *For  $\phi \in C^1$  and  $t \in \mathbb{T}$ ,*

$$\begin{aligned} (\phi, Z_t^K) = & (\Theta_t \phi, Z_0^K) \\ & + \sqrt{K} \int_0^t (-(h_{\tilde{A}_s^K}^K - h_{\tilde{A}_s^\infty}^\infty) \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) (n_{\tilde{A}_s^K}^K - n_{\tilde{A}_s^\infty}^\infty), \tilde{A}_s^\infty) ds \\ & + \int_0^t (-h_{\tilde{A}_s^K}^K \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) n_{\tilde{A}_s^K}^K, Z_s^K) ds + \int_0^t (\Theta_{t-s} \phi, d\tilde{M}_s^K). \end{aligned} \quad (19)$$

The main step in proving tightness is the following bound.

4.3.  $\mathbb{E}[\|Z_t^K\|_{W^{-2}}]$  is bounded by a constant (independent of  $K$  and  $t$ ).

**Proposition 14.**

$$\sup_{t \leq T} \sup_{K \geq 1} \mathbb{E}[\|Z_t^K\|_{W^{-2}}] < \infty.$$

The proof is done using representation (19) with  $\phi \in W^2$ . Each term on the RHS is dealt with separately using successive bounds.

First, we need to overcome the fact that the functions  $\phi$  and  $\theta_t \phi$  are defined on different domains,  $\mathbb{T}^* = [0, T^*]$  and  $[0, T^* - t]$ , respectively. The following lemma constructs an extension of  $\theta_t \phi$  to  $\mathbb{T}^*$  in a way that controls the norm.

**Lemma 15.** *Let  $\phi \in W^j$  for some  $j \in \mathbb{N}$  and  $t \in \mathbb{T}$  be fixed. There exists a function  $\psi : \mathbb{T}^* \rightarrow \mathbb{R}$  such that  $\psi(x) = \phi(x+t)$  for  $x \in [0, T^* - t]$ , and  $\psi \in W^j$  with  $\|\psi\|_{W^j} \leq c\|\phi\|_{W^j}$ , where  $c$  is a constant that depends on  $T^*$  and  $j$ , but independent of  $\phi$ .*

*Proof.* Take  $\psi$  such that  $\psi(x) = \phi(x+t)$  for  $x \in [0, T^* - t]$ , and  $\psi^{(j-1)}(x) = \phi^{(j-1)}(2(T^* - t) - x + t)$  for  $x \in (T^* - t, T^*]$ . That is,  $\psi$  is extended by reflecting the  $(j-1)$ th derivative along  $x = T^* - t$ . Then,  $\psi^{(i)}$  is continuous for  $i = 0, 1, \dots, j-1$ . Note that  $\psi^{(j)}$  does not exist at  $x = T^* - t$ , unless  $\phi^{(j)}(T^*) = 0$ .

It remains to show that  $\|\psi\|_{W^j} \leq c\|\phi\|_{W^j}$ . For  $i = j-1, j$ ,

$$\begin{aligned} \int_{\mathbb{T}^*} (\psi^{(i)}(x))^2 dx &= \int_0^{T^*-t} (\psi^{(i)}(x))^2 dx + \int_{T^*-t}^{T^*} (\psi^{(i)}(x))^2 dx \\ &= \int_t^{T^*} (\phi^{(i)}(x))^2 dx + \int_{T^*-t}^{T^*} (\phi^{(i)}(2(T^* - t) - x + t))^2 dx. \end{aligned}$$

For  $i = 0, 1, \dots, j-2$ ,

$$\begin{aligned} \int_{\mathbb{T}^*} (\psi^{(i)}(x))^2 dx &= \int_0^{T^*-t} (\psi^{(i)}(x))^2 dx + \int_{T^*-t}^{T^*} (\psi^{(i)}(x))^2 dx \\ &= \int_t^{T^*} (\phi^{(i)}(x))^2 dx + \int_{T^*-t}^{T^*} (\psi^{(i)}(x))^2 dx. \end{aligned}$$

For the last integral, note that for  $x \in (T^* - t, T^*]$ ,

$$\begin{aligned} \psi^{(i)}(x) &= \psi^{(i)}(T^* - t) + \int_{T^*-t}^x \psi^{(i+1)}(y) dy \\ &= \phi^{(i)}(T^*) + \int_{T^*-t}^x \psi^{(i+1)}(y) dy, \end{aligned}$$

which can be obtained recursively and be expressed in terms of  $\phi$ . Finally, as  $\phi \in W^j$ , we have  $\phi \in C^{j-1}$  and  $\|\phi\|_{C^{j-1}} = \max_{0 \leq i \leq j-1} \sup_{x \in \mathbb{T}^*} |\phi^{(i)}(x)| < \infty$ . Thus, with  $\|\phi\|_{C^{j-1}} \leq \|\phi\|_{W^j}$  and that  $\mathbb{T}^*$  is a bounded interval, we can bound  $\|\psi\|_{W^j}$  in terms of  $T^*$  and  $\|\phi\|_{W^j}$  and write  $\|\psi\|_{W^j} \leq c\|\phi\|_{W^j}$ .  $\square$

In the sequel,  $\Theta_t \phi$  will refer to its own extension to  $\mathbb{T}^*$ . We immediately get the following inequalities.

$$\|\Theta_t \phi\|_{W^j} \leq c\|\phi\|_{W^j}, \quad \text{and for any } x \in \mathbb{T}^*, |\Theta_t \phi(x)| \leq c\|\phi\|_{W^j}. \quad (20)$$

Next, we give some bounds that are useful in proving Proposition 14.

**Proposition 16.** *Suppose (A2) and (A3) hold. Then we have, for  $t \in \mathbb{T}$  and for all  $x \in \mathbb{T}^*$ ,*

$$\sqrt{K} |h_{\bar{A}_t^K}^K - h_{\bar{A}_t^\infty}^\infty|(x) \leq c(1 + \|Z_t^K\|_{W^{-4}})$$

and

$$\sqrt{K} |n_{\bar{A}_t^K}^K - n_{\bar{A}_t^\infty}^\infty|(x) \leq c(1 + \|Z_t^K\|_{W^{-4}}).$$

*Proof.* We prove only the first inequality, as the second is similar. By the triangle inequality, we have

$$\begin{aligned} |h_{\bar{A}_t^K}^K - h_{\bar{A}_t^\infty}^\infty|(x) &\leq \|h_{\bar{A}_t^K}^K - h_{\bar{A}_t^K}^\infty\|_\infty + \|h_{\bar{A}_t^K}^\infty - h_{\bar{A}_t^\infty}^\infty - \partial_A h_{\bar{A}_t^\infty}^\infty(\bar{A}_t^K - \bar{A}_t^\infty)\|_\infty \\ &\quad + \|\partial_A h_{\bar{A}_t^\infty}^\infty\|_{\mathbb{L}^{-4}} \|\bar{A}_t^K - \bar{A}_t^\infty\|_{W^{-4}}. \end{aligned}$$

Multiplying by  $\sqrt{K}$  and with some manipulation, we have

$$\begin{aligned} \sqrt{K}|h_{\bar{A}_t^K}^K - h_{\bar{A}_t^\infty}^\infty|(x) &\leq \sqrt{K} \sup_A \|h_A^K - h_A^\infty\|_\infty \\ &\quad + \frac{\|Z_t^K\|_{W^{-4}}}{\|\bar{A}_t^K - \bar{A}_t^\infty\|_{W^{-4}}} \|h_{\bar{A}_t^K}^\infty - h_{\bar{A}_t^\infty}^\infty - \partial_A h_{\bar{A}_t^\infty}^\infty (\bar{A}_t^K - \bar{A}_t^\infty)\|_\infty + c_1 \|Z_t^K\|_{W^{-4}}, \end{aligned}$$

where the bound in the last term is due to (A3). It then follows by (A2) and (A3) that  $\sqrt{K}|h_{\bar{A}_t^K}^K - h_{\bar{A}_t^\infty}^\infty|(x) \leq c_2 + c_3 \|Z_t^K\|_{W^{-4}}$ .  $\square$

The following result follows immediately from Proposition 16.

**Proposition 17.** *Suppose (A2) and (A3) hold. For any  $f \in W^j$ ,  $j \in \mathbb{N}$ , and  $t \in \mathbb{T}$ ,*

$$\sup_{x \in \mathbb{T}^*} |\sqrt{K}(L_{\bar{A}_t^K}^K - L_{\bar{A}_t^\infty}^\infty)f|(x) \leq c(1 + \|Z_t^K\|_{W^{-4}})\|f\|_{W^j}.$$

As the operator  $L_A^K$  maps  $W^j$  into  $W^{j-1}$ , we introduce  $\widehat{L}_A^K f = -h_A^K f + f(0)n_A^K$ , so that  $L_A^K f = f' + \widehat{L}_A^K f$ , and let  $\mathcal{L}^{j,k} = L(W^j, W^k)$ , the space of linear operators from  $W^j$  to  $W^k$ .

**Proposition 18.** *Suppose (A2) holds. Then, we have*

$$\sup_{K,A} \|\widehat{L}_A^K\|_{\mathcal{L}^{j,j}} \leq c, \quad j \leq 3; \quad (\text{i})$$

$$\sup_{K,A} \|L_A^K\|_{\mathcal{L}^{j,j-1}} \leq c, \quad 2 \leq j \leq 4. \quad (\text{ii})$$

*Proof.* For  $f \in W^j$ , using triangle inequality and (11),

$$\|\widehat{L}_A^K f\|_{W^j} \leq \|h_A^K\|_{C^j} \|f\|_{W^j} + \|f\|_{W^j} \|n_A^K\|_{W^j} \leq c_1 \|f\|_{W^j}$$

due to embedding and (A2). Thus, (i) follows. For (ii),

$$\|L_A^K f\|_{W^{j-1}} = \|f'\|_{W^{j-1}} + \|\widehat{L}_A^K f\|_{W^{j-1}} \leq \|f\|_{W^j} + c_1 \|f\|_{W^{j-1}} \leq c_2 \|f\|_{W^j},$$

by (i) and embedding. Thus, (ii) follows.  $\square$

Recall also the following bounds, obtained in [7]:

$$(1, \bar{A}_t^\infty) \leq (1, \bar{A}_0^\infty) e^{ct} \quad (21)$$

$$\mathbb{E}[(1, \bar{A}_t^K)] \leq (1, \bar{A}_0^K) e^{ct}. \quad (22)$$

*Completion of proof of Proposition 14.* Let  $\phi \in W^2$ . We bound each term on the RHS of (19), and use repeatedly (20). For the first term,

$$|(\Theta_t \phi, Z_0^K)| \leq \|\Theta_t \phi\|_{W^2} \|Z_0^K\|_{W^{-2}} \leq c_1 \|\phi\|_{W^2} \|Z_0^K\|_{W^{-2}}.$$

For the second term, using Proposition 17,

$$\begin{aligned} &\left| \sqrt{K} \int_0^t ((L_{\bar{A}_s^K}^K - L_{\bar{A}_s^\infty}^\infty) \Theta_{t-s} \phi, \bar{A}_s^\infty) ds \right| \\ &\leq c_2 \int_0^t (1 + \|Z_s^K\|_{W^{-4}}) \|\Theta_{t-s} \phi\|_{W^2} (1, \bar{A}_s^\infty) ds \\ &\leq c_3 \|\phi\|_{W^2} (1, \bar{A}_0^\infty) e^{c_4 t} \int_0^t (1 + \|Z_s^K\|_{W^{-2}}) ds \end{aligned}$$

by (21) and the embedding  $W^{-2} \hookrightarrow W^{-4}$ . For the third term, using Proposition 18(i),

$$\begin{aligned} \left| \int_0^t (-\widehat{L}_{\bar{A}_s^K} \Theta_{t-s} \phi, Z_s^K) ds \right| &\leq c_5 \int_0^t \|\widehat{L}_{\bar{A}_s^K} \Theta_{t-s} \phi\|_{W^2} \|Z_s^K\|_{W^{-2}} ds \\ &\leq c_5 \int_0^t \|\widehat{L}_{\bar{A}_s^K}\|_{\mathcal{L}^{2,2}} \|\Theta_{t-s} \phi\|_{W^2} \|Z_s^K\|_{W^{-2}} ds \\ &\leq c_6 \|\phi\|_{W^2} \int_0^t \|Z_s^K\|_{W^{-2}} ds. \end{aligned}$$

For the forth term, we write  $\int_0^t \Theta_{t-s}^* d\tilde{M}_s^K$  for the map  $f \mapsto \int_0^t (\Theta_{t-s} f, d\tilde{M}_s^K)$ . Then,

$$\left| \int_0^t (\Theta_{t-s} \phi, d\tilde{M}_s^K) \right| \leq \|\phi\|_{W^2} \left\| \int_0^t (\Theta_{t-s}^* d\tilde{M}_s^K) \right\|_{W^{-2}}.$$

Note that  $(\int_0^t (\Theta_{t-s} f, d\tilde{M}_s^K))_{t \in \mathbb{T}}$  is not a martingale, but for each fix  $t$ ,  $(\int_0^r (\Theta_{t-s} f, d\tilde{M}_s^K))_r$  is. Let  $t \in \mathbb{T}$  be fixed. For  $r \leq t$ , by the Riesz Representation Theorem and Parseval's identity,

$$\begin{aligned} \mathbb{E} \left[ \left\| \int_0^r \Theta_{t-s}^* d\tilde{M}_s^K \right\|_{W^{-2}}^2 \right] &= \mathbb{E} \left[ \sum_{l \geq 1} \left( \int_0^r (\Theta_{t-s} p_l^2, d\tilde{M}_s^K) \right)^2 \right] \\ &= \sum_{l \geq 1} \mathbb{E} \left[ \left\langle \int_0^r (\Theta_{t-s} p_l^2, d\tilde{M}_s^K) \right\rangle_r \right] \\ &= \sum_{l \geq 1} \mathbb{E} \left[ \int_0^r \left( (\Theta_{t-s} p_l^2(0))^2 w_{\bar{A}_s^K}^K + h_{\bar{A}_s^K}^K (\Theta_{t-s} p_l^2)^2 \right. \right. \\ &\quad \left. \left. - 2\Theta_{t-s} p_l^2(0) h_{\bar{A}_s^K}^K \widehat{m}_{\bar{A}_s^K}^K \Theta_{t-s} p_l^2, \bar{A}_s^K \right) ds \right]. \end{aligned}$$

It then follows from (12), (C0) and (22) that this quantity is bounded by  $c_7(1, \bar{A}_0^K) e^{c_8 r} r$ . Taking  $r = t$ , we have

$$\mathbb{E} \left[ \left\| \int_0^t \Theta_{t-s}^* d\tilde{M}_s^K \right\|_{W^{-2}}^2 \right] \leq c_7(1, \bar{A}_0^K) e^{c_8 t} t. \quad (23)$$

Now, putting all together with triangle inequality,

$$\begin{aligned} |(\phi, Z_t^K)| &\leq c_9 \left\{ \|\phi\|_{W^2} \|Z_0^K\|_{W^{-2}} + \|\phi\|_{W^2} (1, \bar{A}_0^\infty) e^{c_4 t} \int_0^t (1 + \|Z_s^K\|_{W^{-2}}) ds \right. \\ &\quad \left. + \|\phi\|_{W^2} \int_0^t \|Z_s^K\|_{W^{-2}} ds \right\} + \|\phi\|_{W^2} \left\| \int_0^t \Theta_{t-s}^* d\tilde{M}_s^K \right\|_{W^{-2}}. \end{aligned}$$

This gives a bound on  $\|Z_t^K\|_{W^{-2}}$ . Taking expectation and using (23), we have for  $t \leq T$ ,

$$\begin{aligned} \mathbb{E}[\|Z_t^K\|_{W^{-2}}] &\leq c_{10} \left\{ \|Z_0^K\|_{W^{-2}} + (1, \bar{A}_0^\infty) e^{c_4 t} t \right. \\ &\quad \left. + (1 + (1, \bar{A}_0^\infty) e^{c_4 t}) \int_0^t \mathbb{E}[\|Z_s^K\|_{W^{-2}}] ds + (1, \bar{A}_0^K)^{1/2} e^{c_{11} t} t^{1/2} \right\}. \end{aligned}$$



It follows by Gronwall's inequality that

$$\begin{aligned} \mathbb{E}[\|Z_t^K\|_{W^{-2}}] \\ \leq c_{10}\{\|Z_0^K\|_{W^{-2}} + (1, \bar{A}_0^\infty)e^{c_4 T}T + (1, \bar{A}_0^K)^{1/2}e^{c_{11}T}T^{1/2}\}e^{c_{10}(1+(1, \bar{A}_0^\infty)e^{c_4 T})t}. \end{aligned}$$

Finally, taking supremum over  $t$  and  $K$ , this quantity is finite due to (A4) and (C3).  $\square$

**4.4. Proof of tightness.** It remains to check the tightness condition (T2) only, as (T1) holds by Proposition 14. The conditions (i) and (ii) are verified in a few steps.

Proceeding from Theorem 11, we let

$$\Lambda_t^K f = \sqrt{K}((L_{\bar{A}_t^K}^K - L_{\bar{A}_t^\infty}^\infty)f, \bar{A}_t^\infty) + (L_{\bar{A}_t^K}^K f, Z_t^K)$$

and

$$\Gamma_t^K f = (f^2(0)w_{\bar{A}_t^K}^K + h_{\bar{A}_t^K}^K f^2 - 2f(0)h_{\bar{A}_t^K}^K \hat{m}_{\bar{A}_t^K}^K f, \bar{A}_t^K).$$

**Proposition 19.** *Let  $j \in \{3, 4\}$ . For  $f \in W^j$ ,*

$$|\Lambda_t^K f| \leq c_1 \|f\|_{W^j} (1 + (1, \bar{A}_0^\infty)e^{c_2 t}) (1 + \|Z_t^K\|_{W^{-(j-1)}}).$$

*Proof.* For  $f \in W^j$ , we have  $L_{\bar{A}_t^K}^K f \in W^{j-1}$  and

$$\begin{aligned} |\Lambda_t^K f| &\leq |\sqrt{K}((L_{\bar{A}_t^K}^K - L_{\bar{A}_t^\infty}^\infty)f, \bar{A}_t^\infty)| + |(L_{\bar{A}_t^K}^K f, Z_t^K)| \\ &\leq (|\sqrt{K}(L_{\bar{A}_t^K}^K - L_{\bar{A}_t^\infty}^\infty)f|, \bar{A}_t^\infty) + \|L_{\bar{A}_t^K}^K\|_{\mathcal{L}^{j,j-1}} \|f\|_{W^j} \|Z_t^K\|_{W^{-(j-1)}} \\ &\leq c_1 (1 + \|Z_t^K\|_{W^{-4}}) \|f\|_{W^j} (1, \bar{A}_t^\infty) + c_2 \|f\|_{W^j} \|Z_t^K\|_{W^{-(j-1)}} \end{aligned}$$

due to Propositions 17 and 18. Thus, by (21) and the embedding  $W^{-(j-1)} \hookrightarrow W^{-j}$ , we have

$$|\Lambda_t^K f| \leq c_3 (1 + \|Z_t^K\|_{W^{-(j-1)}}) \|f\|_{W^j} (1, \bar{A}_0^\infty)e^{c_4 t} + c_2 \|f\|_{W^j} \|Z_t^K\|_{W^{-(j-1)}}.$$

The statement now follows.  $\square$

**Proposition 20.**

$$\left| \sum_{l \geq 1} \Gamma_t^K p_l^j \right| \leq c_1 (1, \bar{A}_t^K).$$

*Proof.* This follows directly from (C0) and (12).  $\square$

**Proposition 21.**

$$\sup_{K \geq 1} \mathbb{E} \left[ \sup_{t \leq T} \|Z_t^K\|_{W^{-3}} \right] < \infty.$$

*Proof.* Let  $f \in W^3$ . Using Proposition 19, we have

$$\begin{aligned} |(f, Z_t^K)| &\leq \|f\|_{W^3} \|Z_0^K\|_{W^{-3}} \\ &\quad + c_1 \|f\|_{W^3} (1 + (1, \bar{A}_0^\infty)e^{c_2 t}) \int_0^t (1 + \|Z_s^K\|_{W^{-2}}) ds + \|f\|_{W^3} \|\tilde{M}_t^K\|_{W^{-3}}. \end{aligned}$$

This gives a bound on  $\|Z_t^K\|_{W^{-3}}$  and consequently,

$$\begin{aligned} \sup_{t \leq T} \|Z_t^K\|_{W^{-3}} &\leq \|Z_0^K\|_{W^{-3}} \\ &\quad + c_1(1 + (1, \bar{A}_0^\infty)e^{c_2 T}) \int_0^T (1 + \|Z_s^K\|_{W^{-2}}) ds + \sup_{t \leq T} \|\tilde{M}_t^K\|_{W^{-3}}. \end{aligned} \quad (24)$$

Now, by the Riesz Representation Theorem and Parseval's identity, we have

$$\begin{aligned} \mathbb{E} \left[ \sup_{t \leq T} \|\tilde{M}_t^K\|_{W^{-3}}^2 \right] &= \mathbb{E} \left[ \sup_{t \leq T} \sum_{l \geq 1} (\tilde{M}_t^{p_l^3, K})^2 \right] \\ &\leq \mathbb{E} \left[ \sum_{l \geq 1} \sup_{t \leq T} (\tilde{M}_t^{p_l^3, K})^2 \right] \leq 4 \sum_{l \geq 1} \mathbb{E} \left[ \langle \tilde{M}^{p_l^3, K} \rangle_T \right] \end{aligned}$$

using Doob's inequality. It then follows by Proposition 20 and inequality (22) that

$$\mathbb{E} \left[ \sup_{t \leq T} \|\tilde{M}_t^K\|_{W^{-3}}^2 \right] \leq 4 \sum_{l \geq 1} \mathbb{E} \left[ \int_0^T \Gamma_s^K p_l^3 ds \right] \leq c_5(1, \bar{A}_0^K) e^{c_6 T} T.$$

Therefore, taking expectation in (24), we obtain

$$\begin{aligned} \mathbb{E} \left[ \sup_{t \leq T} \|Z_t^K\|_{W^{-3}} \right] &\leq c_7 \left\{ \|Z_0^K\|_{W^{-3}} \right. \\ &\quad \left. + (1 + (1, \bar{A}_0^\infty)e^{c_2 T}) \int_0^T (1 + \mathbb{E}[\|Z_s^K\|_{W^{-2}}]) ds + (1, \bar{A}_0^K)^{1/2} e^{c_8 T} T^{1/2} \right\}. \end{aligned}$$

Noting that  $\mathbb{E}[\|Z_s^K\|_{W^{-2}}]$  is bounded by Proposition 14, and using (A4) and (C3), complete the proof.  $\square$

**Proposition 22.** *Conditions (i) and (ii) of (T2) hold for  $W^{-4}$ :*

$$\sup_{K \geq 1} \mathbb{E} \left[ \sup_{t \leq T} \|\Lambda_t^K\|_{W^{-4}} \right] \leq c_T \quad (i)$$

$$\sup_{K \geq 1} \mathbb{E} \left[ \sup_{t \leq T} \left| \sum_{l \geq 1} \Gamma_t^K p_l^4 \right| \right] \leq c_T. \quad (ii)$$

*Proof.* From Proposition 19 with  $j = 4$ , we have

$$\|\Lambda_t^K\|_{W^{-4}} \leq c_1(1 + (1, \bar{A}_0^\infty)e^{c_2 t})(1 + \|Z_t^K\|_{W^{-3}}).$$

Taking supremum over  $t \leq T$  and expectation, we have

$$\mathbb{E} \left[ \sup_{t \leq T} \|\Lambda_t^K\|_{W^{-4}} \right] \leq c_1(1 + (1, \bar{A}_0^\infty)e^{c_2 T}) \left( 1 + \mathbb{E} \left[ \sup_{t \leq T} \|Z_t^K\|_{W^{-3}} \right] \right),$$

which is bounded in  $K$  by Proposition 21. Thus, condition (i) holds.

Now we verify condition (ii). From Proposition 20,

$$\mathbb{E} \left[ \sup_{t \leq T} \left| \sum_{l \geq 1} \Gamma_t^K p_l^4 \right| \right] \leq c_3 \mathbb{E} \left[ \sup_{t \leq T} (1, \bar{A}_t^K) \right].$$

But,

$$(1, \bar{A}_t^K) \leq (1, \bar{A}_0^K) + c_4 \int_0^t (1, \bar{A}_s^K) ds + \frac{1}{\sqrt{K}} \tilde{M}_t^{1, K}$$

and for  $S \leq T$ ,

$$\mathbb{E} \left[ \sup_{t \leq S} (1, \bar{A}_t^K) \right] \leq (1, \bar{A}_0^K) + c_4 \int_0^S \mathbb{E} \left[ \sup_{u \leq s} (1, \bar{A}_u^K) \right] ds + \frac{1}{\sqrt{K}} \mathbb{E} \left[ \sup_{t \leq T} \tilde{M}_t^{1,K} \right].$$

It follows by Gronwall's inequality that

$$\mathbb{E} \left[ \sup_{t \leq T} (1, \bar{A}_t^K) \right] \leq \left\{ (1, \bar{A}_0^K) + \frac{1}{\sqrt{K}} \mathbb{E} \left[ \sup_{t \leq T} \tilde{M}_t^{1,K} \right] \right\} e^{c_4 T},$$

where by Doob's inequality,

$$\mathbb{E} \left[ \sup_{t \leq T} \tilde{M}_t^{1,K} \right]^2 \leq \mathbb{E} \left[ \sup_{t \leq T} (\tilde{M}_t^{1,K})^2 \right] \leq 4 \mathbb{E} [\langle \tilde{M}^{1,K} \rangle_T] \leq c_5 (1, \bar{A}_0^K) e^{c_6 T}.$$

Therefore, condition (ii) follows, using (C3).  $\square$

**Corollary 23.** *Both sequences  $Z^K$  and  $\tilde{M}^K$  are tight in  $\mathbb{D}(\mathbb{T}, W^{-4})$ .*

**4.5. C-tightness of  $Z^K$  and  $\tilde{M}^K$ .** We can further show that  $Z^K$  and  $\tilde{M}^K$  are C-tight, that is, the two sequences are tight and all limit points of the sequences are continuous.

**Proposition 24.** *The sequence  $Z^K$  is C-tight and all limit points of  $Z^K$  are elements of  $\mathbb{C}(\mathbb{T}, W^{-4})$ .*

*Proof.* We have established that  $Z^K$  is tight, it remains to show that (see e.g. [9, Proposition VI 3.26(iii)]), for all  $u \in \mathbb{T}$  and  $\epsilon > 0$ ,

$$\lim_{K \rightarrow \infty} \mathbb{P} \left( \sup_{t \leq u} \|\Delta Z_t^K\|_{W^{-4}} > \epsilon \right) = 0.$$

We observe that  $Z^K$  jumps when  $A^K$  jumps, which occurs when there is a birth or a death. Thus, for  $f \in W^4$ , we have

$$\begin{aligned} |(f, \Delta Z_t^K)| &= |(f, Z_t^K - Z_{t-}^K)| = \frac{1}{\sqrt{K}} |(f, A_t^K - A_{t-}^K)| \\ &\leq \frac{1}{\sqrt{K}} \max \left\{ \sup_{x \in \mathbb{T}^*} \check{\xi}_{A_t^K}^K(x) |f(0)|, \sup_{x \in \mathbb{T}^*} |\hat{\xi}_{A_t^K}^K(x) f(0) - f(x)| \right\} \\ &\leq \frac{1}{\sqrt{K}} \|f\|_{W^4} (1 + \Xi) \end{aligned}$$

by (A1), giving  $\|\Delta Z_t^K\|_{W^{-4}} \leq \frac{1}{\sqrt{K}} (1 + \Xi)$ . Hence,

$$\mathbb{P} \left( \sup_{t \leq u} \|\Delta Z_t^K\|_{W^{-4}} > \epsilon \right) \leq \frac{1}{\epsilon} \mathbb{E} \left[ \sup_{t \leq u} \|\Delta Z_t^K\|_{W^{-4}} \right] \leq \frac{1}{\epsilon} \frac{1}{\sqrt{K}} (1 + \mathbb{E}[\Xi]),$$

which converges to zero as  $K$  tends to infinity.  $\square$

**Corollary 25.** *The sequence of martingales  $\tilde{M}^K$  is C-tight and all limit points of  $\tilde{M}^K$  are elements of  $\mathbb{C}(\mathbb{T}, W^{-4})$ .*

*Proof.* As  $Z^K$  and  $\tilde{M}^K$  have the same discontinuities,  $\Delta Z_t^K = \Delta \tilde{M}_t^K$  and it follows that  $\tilde{M}^K$  satisfies the conditions of being C-tight.  $\square$

#### 4.6. Convergence of $\tilde{M}^K$ and $Z^K$ .

**Proposition 26.** *The sequence  $\tilde{M}^K$  converges weakly to  $\tilde{M}^\infty$  such that for any  $f \in W^4$ ,  $\tilde{M}^{f,\infty} \equiv (f, \tilde{M}^\infty)$ ,  $t \in \mathbb{T}$ , is a continuous Gaussian martingale with predictable quadratic variation*

$$\langle \tilde{M}^{f,\infty} \rangle_t = \int_0^t (f^2(0)w_{\tilde{A}_s^\infty}^\infty + h_{\tilde{A}_s^\infty}^\infty f^2 - 2f(0)h_{\tilde{A}_s^\infty}^\infty \hat{m}_{\tilde{A}_s^\infty}^\infty f, \bar{A}_s^\infty) ds. \quad (25)$$

*Proof.* Let  $f \in W^4$ . Recall from the proof of Proposition 24 that

$$\sup_{s \leq t} |\Delta \tilde{M}_s^{f,K}|^2 = \sup_{s \leq t} |\Delta(f, Z_s^K)|^2 \leq \frac{1}{K} \|f\|_{W^4}^2 (1 + \Xi)^2.$$

Thus,

$$\sup_{K \geq 1} \mathbb{E} \left[ \sup_{s \leq t} |\Delta \tilde{M}_s^{f,K}|^2 \right] \leq \sup_{K \geq 1} \mathbb{E} \left[ \frac{1}{K} \|f\|_{W^4}^2 (1 + \Xi)^2 \right],$$

which is finite by (A1). Therefore,  $\sup_{s \leq t} |\Delta \tilde{M}_s^{f,K}|$  is uniformly integrable and converges to zero in probability for all  $t \in \mathbb{T}$ . All limit points of  $\tilde{M}^{f,K}$  are continuous (from Corollary 25) and  $\langle \tilde{M}^{f,K} \rangle_t$  converges to (25). By [9, Theorem VIII 3.12(iv)]  $\tilde{M}^{f,K}$  converges to a continuous martingale  $\tilde{M}^{f,\infty}$  with predictable quadratic variation in (25). The limiting process is Gaussian as the predictable quadratic variation is deterministic.

Tightness of  $\tilde{M}^K$  implies that there exists a subsequence that converges. Suppose  $M$  and  $N$  both are accumulation points of  $\tilde{M}^K$ . Then, we have  $(f, M) = \tilde{M}^{f,\infty} = (f, N)$  for every  $f \in W^4$ , and thus, we must have  $M = N$  in  $W^{-4}$ . Therefore, we can conclude that  $\tilde{M}^K$  converges to  $\tilde{M}^\infty$ , where  $\tilde{M}^\infty$  is defined such that  $(f, \tilde{M}^\infty) = \tilde{M}^{f,\infty}$  for every  $f \in W^4$ .  $\square$

**Proposition 27.** *Every limit point  $Z$  of the sequence  $Z^K$  satisfies, for  $\phi \in W^4$  and  $t \in \mathbb{T}$ ,*

$$\begin{aligned} (\phi, Z_t) &= (\Theta_t \phi, Z_0) + \int_0^t (-\partial_A h_{\tilde{A}_s^\infty}^\infty(Z_s) \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) \partial_A n_{\tilde{A}_s^\infty}^\infty(Z_s), \bar{A}_s^\infty) ds \\ &\quad + \int_0^t (-h_{\tilde{A}_s^\infty}^\infty \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) n_{\tilde{A}_s^\infty}^\infty, Z_s) ds + \int_0^t (\Theta_{t-s} \phi, d\tilde{M}_s^\infty). \end{aligned} \quad (26)$$

*Proof.* First, we show that  $\sqrt{K}(h_{\tilde{A}_s^K}^K - h_{\tilde{A}_s^\infty}^\infty)$  converges to  $\partial_A h_{\tilde{A}_s^\infty}^\infty(Z_s)$ :

$$\begin{aligned} &|\sqrt{K}(h_{\tilde{A}_s^K}^K - h_{\tilde{A}_s^\infty}^\infty) - \partial_A h_{\tilde{A}_s^\infty}^\infty(Z_s)| \\ &\leq \sqrt{K}|h_{\tilde{A}_s^K}^K - h_{\tilde{A}_s^\infty}^\infty| + \sqrt{K}|h_{\tilde{A}_s^\infty}^\infty - h_{\tilde{A}_s^\infty}^\infty - \partial_A h_{\tilde{A}_s^\infty}^\infty(\bar{A}_s^K - \bar{A}_s^\infty)| \\ &\quad + |\partial_A h_{\tilde{A}_s^\infty}^\infty(Z_s^K) - \partial_A h_{\tilde{A}_s^\infty}^\infty(Z_s)| \\ &\leq \sqrt{K} \sup_A \|h_A^K - h_A^\infty\|_\infty \\ &\quad + \frac{\|Z_s^K\|_{W^{-4}}}{\|\bar{A}_s^K - \bar{A}_s^\infty\|_{W^{-4}}} \|h_{\tilde{A}_s^\infty}^\infty - h_{\tilde{A}_s^\infty}^\infty - \partial_A h_{\tilde{A}_s^\infty}^\infty(\bar{A}_s^K - \bar{A}_s^\infty)\|_\infty \\ &\quad + \|\partial_A h_{\tilde{A}_s^\infty}^\infty\|_{\mathbb{L}^{-4}} \|Z_s^K - Z_s\|_{W^{-4}}, \end{aligned}$$

which converges to zero as  $K$  tends to infinity; the first term by (A2), the second by the definition of Fréchet derivative (A3), and the last term due to  $Z$  being a limit. Similarly,  $\sqrt{K}(n_{\bar{A}_s^K}^K - n_{\bar{A}_s^\infty}^\infty)$  converges to  $\partial_A n_{\bar{A}_s^\infty}^\infty(Z_s)$ . Thus,

$$\begin{aligned} \sqrt{K} \int_0^t & (- (h_{\bar{A}_s^K}^K - h_{\bar{A}_s^\infty}^\infty) \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) (n_{\bar{A}_s^K}^K - n_{\bar{A}_s^\infty}^\infty), \bar{A}_s^\infty) ds \\ & \xrightarrow{K \rightarrow \infty} \int_0^t (- \partial_A h_{\bar{A}_s^\infty}^\infty(Z_s) \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) \partial_A n_{\bar{A}_s^\infty}^\infty(Z_s), \bar{A}_s^\infty) ds \end{aligned}$$

by dominated convergence theorem.

Next, we show that  $\int_0^t (-h_{\bar{A}_s^K}^K \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) n_{\bar{A}_s^K}^K, Z_s^K) ds$  converges to  $\int_0^t (-h_{\bar{A}_s^\infty}^\infty \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) n_{\bar{A}_s^\infty}^\infty, Z_s) ds$ . Using a similar argument as for Proposition 18(i), with (A2),

$$\begin{aligned} & \left| \int_0^t (-h_{\bar{A}_s^K}^K \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) n_{\bar{A}_s^K}^K, Z_s^K) ds \right. \\ & \quad \left. - \int_0^t (-h_{\bar{A}_s^\infty}^\infty \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) n_{\bar{A}_s^\infty}^\infty, Z_s) ds \right| \\ & \leq \int_0^t \| (-h_{\bar{A}_s^K}^K - h_{\bar{A}_s^\infty}^\infty) \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) (n_{\bar{A}_s^K}^K - n_{\bar{A}_s^\infty}^\infty) \|_{W^4} \|Z_s^K\|_{W^{-4}} ds \\ & \quad + \int_0^t \| -h_{\bar{A}_s^\infty}^\infty \Theta_{t-s} \phi + \Theta_{t-s} \phi(0) n_{\bar{A}_s^\infty}^\infty \|_{W^4} \|Z_s^K - Z_s\|_{W^{-4}} ds \\ & \leq \int_0^t c_1 \|\phi\|_{W^4} (\|h_{\bar{A}_s^K}^K - h_{\bar{A}_s^\infty}^\infty\|_{C^4} + \|n_{\bar{A}_s^K}^K - n_{\bar{A}_s^\infty}^\infty\|_{W^4}) \|Z_s^K\|_{W^{-4}} ds \\ & \quad + \int_0^t c_2 \|\phi\|_{W^4} (\|h_{\bar{A}_s^\infty}^\infty\|_{C^4} + \|n_{\bar{A}_s^\infty}^\infty\|_{W^4}) \|Z_s^K - Z_s\|_{W^{-4}} ds, \end{aligned}$$

which converges to 0 as  $K \rightarrow \infty$ .

Together with the convergence of  $Z_0^K$  in (A4) and the convergence of  $\tilde{M}^K$  established in Proposition 26, the proof is complete.  $\square$

It remains to show the uniqueness of the solution to Equation (26).

**Proposition 28.** *Suppose that  $Z$  and  $Y$  both are solutions to Equation (26) in Proposition 27 with  $Z_0 = Y_0$ , then  $Z = Y$ .*

*Proof.* First, note that Proposition 18(i) remains true if we replace  $\hat{L}_A^K$  with  $\hat{L}_A^\infty$ :  $f \mapsto -h_A^\infty f + f(0)n_A^\infty$ , for  $j \leq 4$ , due to (A2). Now, let  $\phi \in W^4$  and  $t \in \mathbb{T}$ , by

triangle inequality, we have

$$\begin{aligned}
& |(\phi, Z_t - Y_t)| \\
& \leq \int_0^t (|\partial_A h_{\bar{A}_s^\infty}^\infty(Z_s - Y_s)| |\Theta_{t-s}\phi| + |\Theta_{t-s}\phi(0)| |\partial_A n_{\bar{A}_s^\infty}^\infty(Z_s - Y_s)|, \bar{A}_s^\infty) ds \\
& \quad + \int_0^t \| -h_{\bar{A}_s^\infty}^\infty \Theta_{t-s}\phi + \Theta_{t-s}\phi(0) n_{\bar{A}_s^\infty}^\infty \|_{W^4} \|Z_s - Y_s\|_{W^{-4}} ds \\
& \leq \int_0^t c_1 \|\phi\|_{W^4} (\|\partial_A h_{\bar{A}_s^\infty}^\infty\|_{\mathbb{L}^{-4}} + \|\partial_A n_{\bar{A}_s^\infty}^\infty\|_{\mathbb{L}^{-4}}) \|Z_s - Y_s\|_{W^{-4}} (1, \bar{A}_s^\infty) ds \\
& \quad + \int_0^t \|\widehat{L}_{\bar{A}_s^\infty}^\infty\|_{\mathcal{L}^{4,4}} \|\Theta_{t-s}\phi\|_{W^4} \|Z_s - Y_s\|_{W^{-4}} ds \\
& \leq c_2 \|\phi\|_{W^4} (1 + (1, \bar{A}_0^\infty) e^{c_3 T}) \int_0^t \|Z_s - Y_s\|_{W^{-4}} ds.
\end{aligned}$$

It then follows by Gronwall's inequality that  $\|Z_t - Y_t\|_{W^{-4}} = 0$ . Therefore,  $Z = Y$ .  $\square$

Lastly, we note that Equation (26) is the same as Equation (13). This is straightforward and the proof is omitted.

**Proposition 29.** *The limiting process  $Z$  satisfies Equation (13), for any  $f \in W^4$  and  $t \in \mathbb{T}$ .*

## 5. PROOFS OF COROLLARY 9 AND PROPOSITION 10

*Proof of Corollary 9.* The SPDE representation follows by direct calculation. To show that  $(\tilde{M}^\infty)_{t,f}$  is Gaussian, we use the Cramér-Wold device, by showing that for all  $f_1, \dots, f_n$  in  $W^4$ ,  $((\tilde{M}_t^{f_1, \infty}, \dots, \tilde{M}_t^{f_n, \infty}))_{t \geq 0}$  is Gaussian. This is equivalent to showing that for all  $\alpha_1, \dots, \alpha_n \geq 0$ ,  $(\alpha_1 \tilde{M}_t^{f_1, \infty} + \dots + \alpha_n \tilde{M}_t^{f_n, \infty})_{t \in \mathbb{T}}$  is Gaussian, which is true observing that

$$\alpha_1 \tilde{M}_t^{f_1, \infty} + \dots + \alpha_n \tilde{M}_t^{f_n, \infty} = \tilde{M}_t^{(\alpha_1 f_1 + \dots + \alpha_n f_n), \infty}.$$

$\square$

*Proof of Proposition 10.* To prove this, we use representation (26) and obtain<sup>4</sup>

$$\begin{aligned}
\mathbb{E}[(\phi, Z_t)] &= (\Theta_t \phi, Z_0) \\
&+ \int_0^t ( -\mathbb{E}[(g^h(\bar{A}_s^\infty, *, \cdot), Z_s).] \Theta_{t-s}\phi + \phi(t-s) \mathbb{E}[(g^n(\bar{A}_s^\infty, *, \cdot), Z_s).], \bar{A}_s^\infty)_* ds \\
&\quad + \int_0^t \mathbb{E}[( -h_{\bar{A}_s^\infty}^\infty \Theta_{t-s}\phi + \phi(t-s) n_{\bar{A}_s^\infty}^\infty, Z_s)] ds
\end{aligned}$$

---

<sup>4</sup>We write  $((f(*, \cdot), \mu), \nu)_*$  to mean  $\int \int f(x, y) \mu(dy) \nu(dx) = (f, \nu \otimes \mu)$ .

as  $\mathbb{E}[\int_0^t (\Theta_{t-s}\phi, d\tilde{M}_s^\infty) = 0]$ . Defining  $\nu_t : f \mapsto \mathbb{E}[(f, Z_t)]$ , we have

$$\begin{aligned} (\phi, \nu_t) &= (\Theta_t\phi, \nu_0) \\ &+ \int_0^t \left( -(g^h(\bar{A}_s^\infty, *, \cdot), \nu_s) \cdot \Theta_{t-s}\phi + \phi(t-s)(g^n(\bar{A}_s^\infty, *, \cdot), \nu_s) \cdot \bar{A}_s^\infty \right) ds \\ &+ \int_0^t \left( -h_{\bar{A}_s^\infty}^\infty \Theta_{t-s}\phi + \phi(t-s)n_{\bar{A}_s^\infty}^\infty, \nu_s \right) ds \end{aligned} \quad (27)$$

for  $\phi \in W^4$ . Using (20), (21) and (A2), we have

$$\begin{aligned} |(\phi, \nu_t)| &\leq \|\phi\|_{W^4} \|\nu_0\|_{W^{-4}} \\ &+ c_1 \|\phi\|_{W^4} \left( \sup_{A,x} \|g^h(A, x, \cdot)\|_{W^4} + \sup_{A,x} \|g^n(A, x, \cdot)\|_{W^4} \right) (1, \bar{A}_0^\infty) e^{c_2 t} \int_0^t \|\nu_s\|_{W^{-4}} ds \\ &+ c_3 \|\phi\|_{W^4} \int_0^t \|\nu_s\|_{W^{-4}} ds, \end{aligned}$$

which gives, by Gronwall's inequality,

$$\|\nu_t\|_{W^{-4}} \leq c_T \|\nu_0\|_{W^{-4}}.$$

Now, let  $(\phi_k)_k$  be a sequence of functions in  $C^\infty$  that converges to  $\phi \in C^0$ . By dominated convergence theorem, (27) holds for  $\phi \in C^0$ . Moreover,  $\nu : C^0 \rightarrow \mathbb{R}$  is a bounded linear operator. Therefore,  $\nu_t$  can be seen as an element in  $C^{-0}$ , that is, it is a measure.  $\square$

## 6. EXAMPLE: PARAMETERS THAT ARE ESSENTIALLY LINEAR

Suppose the reproduction parameters are of the form  $q_A^K(x) = q(x, |\bar{A}^K|, (g(x, \cdot), \bar{A}^K))$ ,  $q = b, h, m, v$ , where  $g : \mathbb{T}^* \times \mathbb{T}^* \rightarrow \mathbb{R}$  and  $q : \mathbb{T}^* \times \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$ . Suppose that:

- (1) The function  $g$  is element of  $C^{4,4}$ .
- (2) The functions  $b, h, m$  are elements of  $C^{4,1,4}$ ; and for  $q = b, h, m$ ,
  - (a)  $\sup_{x,y,z} |\partial_2 q(x, y, z)| < \infty$
  - (b)  $\sup_{x,y,z} (1+y)^k |\partial_3^k \partial_1^j q(x, y, z)| < \infty$  for  $j = 0, 1, \dots, 4-k$  and  $k = 0, 1, 2, 3, 4$
 where  $\partial_i^j$  denotes the  $j$ th order partial derivative with respect to the  $i$ th variable.
- (3) The function  $v$  is bounded and Lipschitz in the second and the third variables, uniformly in the first variable, i.e.

$$\sup_x |v(x, y_1, z_1) - v(x, y_2, z_2)| \leq c(|y_1 - y_2| + |z_1 - z_2|).$$

Then, together with assumptions (C3), (A1) and (A4), LLN and CLT hold with  $q_A^\infty(x) = q(x, |A|, (g(x, \cdot), A))$  and

$$\begin{aligned} \partial_A q_{A_0}^\infty(B)(x) &= \partial_2 q(x, |A_0|, (g(x, \cdot), A_0))(1, B) \\ &+ \partial_3 q(x, |A_0|, (g(x, \cdot), A_0))(g(x, \cdot), B). \end{aligned}$$

It also follows from Proposition 10 that  $\nu_t : f \mapsto \mathbb{E}[(f, Z_t)]$  is a measure and satisfies the following equation:

$$\begin{aligned} (f, \nu_t) &= (f, \nu_0) + \int_0^t \left\{ (1, \nu_s) \left( f(0) \partial_2 n(*, |\bar{A}_s^\infty|, (g(*, \cdot), \bar{A}_s^\infty).) \right. \right. \\ &\quad \left. \left. - \partial_2 h(*, |\bar{A}_s^\infty|, (g(*, \cdot), \bar{A}_s^\infty).) f, \bar{A}_s^\infty \right)_* \right. \\ &\quad \left. + \left( (f(0) \partial_3 n(*, |\bar{A}_s^\infty|, (g(*, \cdot), \bar{A}_s^\infty).) \right. \right. \\ &\quad \left. \left. - \partial_3 h(*, |\bar{A}_s^\infty|, (g(*, \cdot), \bar{A}_s^\infty).) f) (g(*, \cdot), \nu_s), \bar{A}_s^\infty \right)_* \right\} ds \\ &\quad + \int_0^t \left( f' - h(*, |\bar{A}_s^\infty|, (g(*, \cdot), \bar{A}_s^\infty).) f + f(0) n(*, |\bar{A}_s^\infty|, (g(*, \cdot), \bar{A}_s^\infty).), \nu_s \right)_* ds. \end{aligned}$$

In what follows, we consider a few special cases. We will also see that when  $q_A^K(x)$  is a function of  $|\bar{A}^K|$  only, or is a constant, an explicit expression for the density of the measure  $\mathbb{E}[Z_t]$  can be computed.

**6.1. Special case.** Suppose that the reproduction parameters are of the form  $\bar{q}(x, \frac{(g(x, \cdot), \bar{A}^K)}{1 + |\bar{A}^K|})$ , where  $\bar{q} : \mathbb{T}^* \times \mathbb{R} \rightarrow \mathbb{R}$  bounded and  $g \in C^{4,4}$ . That is, we take  $q(x, y, z) = \bar{q}(x, \frac{z}{1+y})$ . Conditions (2) and (3) on  $q$  above then reduce to  $\bar{q} \in C^{4,4}$  with

- (a)  $\sup_{x,u} |u \partial_2 \bar{q}(x, u)| < \infty$ , for  $\bar{q} = b, h, m, v$
- (b)  $\sup_{x,u} |\partial_1^j \partial_2^k \bar{q}(x, u)| < \infty$ , for  $j = 0, 1, \dots, 4-k$ ,  $k = 0, 1, 2, 3, 4$  and  $\bar{q} = b, h, m$ .

Note that (a) implies the Lipschitz condition. In this case,

$$\partial_A \bar{q}_{A_0}^\infty(B)(x) = \partial_2 \bar{q}\left(x, \frac{(g(x, \cdot), A_0)}{1 + |A_0|}\right) \frac{(1 + |A_0|)(g(x, \cdot), B) - |B|(g(x, \cdot), A_0)}{(1 + |A_0|)^2}.$$

The measure  $\nu_t : f \mapsto \mathbb{E}[(f, Z_t)]$  satisfies the following equation:

$$\begin{aligned} (f, \nu_t) &= (f, \nu_0) + \int_0^t \left( \frac{(1 + |\bar{A}_s^\infty|)(g(*, \cdot), \nu_s) - |\nu_s|(g(*, \cdot), \bar{A}_s^\infty)}{(1 + |\bar{A}_s^\infty|)^2} \right. \\ &\quad \times \left( f(0) \partial_2 n\left(*, \frac{(g(*, \cdot), \bar{A}_s^\infty)}{1 + |\bar{A}_s^\infty|}\right) - \partial_2 h\left(*, \frac{(g(*, \cdot), \bar{A}_s^\infty)}{1 + |\bar{A}_s^\infty|}\right) f, \bar{A}_s^\infty \right)_* ds \\ &\quad \left. + \int_0^t \left( f' - h\left(*, \frac{(g(*, \cdot), \bar{A}_s^\infty)}{1 + |\bar{A}_s^\infty|}\right) f + f(0) n\left(*, \frac{(g(*, \cdot), \bar{A}_s^\infty)}{1 + |\bar{A}_s^\infty|}\right), \nu_s \right)_* ds. \right. \end{aligned}$$

**6.2. Age-and-density-dependent case.** Suppose that the reproduction parameters are of the form  $\tilde{q}(x, |\bar{A}^K|)$ , where  $\tilde{q} : \mathbb{T}^* \times \mathbb{R}_+ \rightarrow \mathbb{R}$  bounded. That is, we take  $q(x, y, z) = \tilde{q}(x, y)$ . Then conditions on  $q$  reduce to  $\tilde{q} \in C^{4,1}$  with

- (a)  $\sup_{x,y} |\partial_2 \tilde{q}(x, y)| < \infty$ , for  $\tilde{q} = b, h, m, v$
- (b)  $\sup_{x,y} |\partial_1^k \tilde{q}(x, y)| < \infty$ , for  $k = 0, 1, 2, 3, 4$  and  $\tilde{q} = b, h, m$ ,

and we have

$$\partial_A \tilde{q}_{A_0}^\infty(B)(x) = \partial_2 \tilde{q}(x, |A_0|) |B|.$$



The measure  $\nu_t : f \mapsto \mathbb{E}[(f, Z_t)]$  satisfies the following equation:

$$\begin{aligned} (f, \nu_t) &= (f, \nu_0) + \int_0^t (1, \nu_s) (f(0) \partial_2 n(\cdot, |\bar{A}_s^\infty|) - \partial_2 h(\cdot, |\bar{A}_s^\infty|) f), \bar{A}_s^\infty) ds \\ &\quad + \int_0^t (f' - h(\cdot, |\bar{A}_s^\infty|) f + f(0) n(\cdot, |\bar{A}_s^\infty|), \nu_s) ds. \end{aligned}$$

**6.3. Density-dependent case.** Suppose that the reproduction parameters are of the form  $\hat{q}(|\bar{A}^K|)$ , where  $\hat{q} : \mathbb{R}_+ \rightarrow \mathbb{R}$ . That is,  $\bar{q}$  is independent of  $x$ . Then the conditions on  $q$  further reduce to  $\hat{q} \in C_b^1$ . In this case,  $\partial_A \hat{q}_{A_0}^\infty(B)(x) = \hat{q}'(|A_0|)|B|$ , and the measure  $\mathbb{E}[Z_t]$  has a density if  $\mathbb{E}[Z_0]$  does.

With  $X_t = |\bar{A}_t^\infty|$ , we can write

$$\begin{aligned} (f, Z_t) &= (f, Z_0) + \int_0^t (-h'(X_s)|Z_s|f + f(0)n'(X_s)|Z_s|, \bar{A}_s^\infty) ds \\ &\quad + \int_0^t (f' - h(X_s)f + f(0)n(X_s), Z_s) ds + \tilde{M}_t^{f, \infty}. \end{aligned}$$

Taking  $f_\lambda(x) = e^{\lambda x}$  and writing  $\tilde{M}_t^\lambda$  for the martingale, we have

$$\begin{aligned} (f_\lambda, Z_t) &= (f_\lambda, Z_0) + \\ &\quad \int_0^t (n(X_s) + n'(X_s)X_s - h'(X_s)(f_\lambda, \bar{A}_s^\infty))(f_0, Z_s) ds \\ &\quad + \int_0^t (\lambda - h(X_s))(f_\lambda, Z_s) ds + \tilde{M}_t^\lambda. \end{aligned} \quad (28)$$

Taking expectation and letting  $\phi(s, \lambda) = n(X_s) + n'(X_s)X_s - h'(X_s)(f_\lambda, \bar{A}_s^\infty)$  and  $\psi(s, \lambda) = \lambda - h(X_s)$ ,

$$\mathbb{E}[(f_\lambda, Z_t)] = \mathbb{E}[(f_\lambda, Z_0)] + \int_0^t \phi(s, \lambda) \mathbb{E}[(f_0, Z_s)] ds + \int_0^t \psi(s, \lambda) \mathbb{E}[(f_\lambda, Z_s)] ds.$$

Solving this gives

$$\begin{aligned} \mathbb{E}[(f_\lambda, Z_t)] &= e^{\int_0^t \psi(s, \lambda) ds} \left\{ \mathbb{E}[(f_\lambda, Z_0)] \right. \\ &\quad \left. + \mathbb{E}[(f_0, Z_0)] \int_0^t \phi(s, \lambda) e^{\int_0^s (\psi(r, 0) - \psi(r, \lambda) + \phi(r, 0)) dr} ds \right\}, \end{aligned}$$

which reduces to

$$\begin{aligned} \mathbb{E}[(f_\lambda, Z_t)] &= e^{\lambda t - \int_0^t h(X_s) ds} \left\{ \mathbb{E}[(f_\lambda, Z_0)] \right. \\ &\quad \left. + \mathbb{E}[(f_0, Z_0)] \int_0^t (n(X_s) + n'(X_s)X_s - h'(X_s)(f_\lambda, \bar{A}_s^\infty)) \right. \\ &\quad \left. \times e^{-\lambda s + \int_0^s (n(X_r) + (n'(X_r) - h'(X_r))X_r) dr} ds \right\}. \end{aligned}$$

Inverting the transform, we obtain an expression for the density. Suppose that  $\mathbb{E}[Z_0]$  has density  $\mathfrak{z}_0(x)$ , then  $\mathbb{E}[Z_t]$  has density  $\mathfrak{z}_t(x)$  and

$$\begin{aligned} \mathfrak{z}_t(x) = e^{-\int_0^t h(X_s)ds} & \left\{ \mathfrak{z}_0(x-t) \mathbf{1}_{x>t} + \mathbb{E}[(1, Z_0)] \left\{ \right. \right. \\ & \left. \left( n(X_{t-x}) + n'(X_{t-x})X_{t-x} \right) e^{\int_0^{t-x} (n(X_r) + (n'(X_r) - h'(X_r))X_r)dr} \mathbf{1}_{x \leq t} \right. \\ & \left. \left. - \int_{(t-x) \vee 0}^t h'(X_s) e^{\int_0^s (n(X_r) + (n'(X_r) - h'(X_r))X_r)dr} a(x-t+s, t) ds \right\} \right\} \end{aligned}$$

where  $a(x, t)$  is the density of  $\bar{A}_t^\infty$ .

We can also write the SPDE of  $Z$ :

$$dZ_t = \left( (n(X_t) + n'(X_t)X_t)|Z_t|\delta_0 - h(X_t)Z_t - h'(X_t)|Z_t|\bar{A}_t^\infty - (Z_t)' \right) dt + d\tilde{M}_t^\infty$$

with

$$\begin{aligned} \langle \tilde{M}^{f, \infty} \rangle_t &= \int_0^t (f^2(0)w(X_t) + h(X_t)f^2 - 2f(0)h(X_t)\hat{m}(X_t)f, \bar{A}_s^\infty)ds \\ &= f^2(0)w(X_t) \int_0^t X_s ds + h(X_t) \int_0^t (f^2, \bar{A}_s^\infty)ds \\ &\quad - 2f(0)h(X_t)\hat{m}(X_t) \int_0^t (f, \bar{A}_s^\infty)ds, \end{aligned}$$

and  $(1, \bar{A}_s^\infty) = X_s = X_0 e^{\int_0^s (n(X_r) - h(X_r))dr}$ .

In fact, we can solve (28) and obtain

$$(f_\lambda, Z_t) = e^{\int_0^t (\lambda - h(X_s))ds} (f_\lambda, Z_0) + \int_0^t e^{\int_s^t (\lambda - h(X_r))dr} \left( \phi(s, \lambda) |Z_s| ds + d\tilde{M}_s^\lambda \right).$$

But,

$$|Z_t| = e^{\int_0^t \varphi(s)ds} |Z_0| + \int_0^t e^{\int_s^t \varphi(r)dr} d\tilde{M}_s^0$$

with  $\varphi(s) = n(X_s) - h(X_s) + (n'(X_s) - h'(X_s))X_s$ . Thus,

$$\begin{aligned} (f_\lambda, Z_t) &= e^{\int_0^t (\lambda - h(X_s))ds} (f_\lambda, Z_0) \\ &\quad + \int_0^t e^{\int_s^t (\lambda - h(X_r))dr} \left( \phi(s, \lambda) \left( e^{\int_0^s \varphi(r)dr} |Z_0| + \int_0^s e^{\int_u^s \varphi(r)dr} d\tilde{M}_u^0 \right) ds + d\tilde{M}_s^\lambda \right) \end{aligned}$$

with

$$\langle \tilde{M}^0, \tilde{M}^\lambda \rangle_t = \int_0^t \left( w(X_s)X_s - h(X_s)\hat{m}(X_s)X_s + h(X_s)(1 - \hat{m}(X_s))(f_\lambda, \bar{A}_s^\infty) \right) ds.$$

**6.4. Classical case.** Assume constant parameters  $b, h, m$  and  $v$ , then we have for a test function  $f$ ,

$$(f, Z_t) = (f, Z_0) + \int_0^t (f' - hf + f(0)n, Z_s)ds + \tilde{M}_t^{f, \infty}.$$

Taking  $f_\lambda(x) = e^{\lambda x}$  and writing  $\tilde{M}_t^\lambda$  for the martingale, we have

$$(f_\lambda, Z_t) = (f_\lambda, Z_0) + (\lambda - h) \int_0^t (f_\lambda, Z_s)ds + n \int_0^t (1, Z_s)ds + \tilde{M}_t^\lambda \quad (29)$$

with

$$\langle \tilde{M}^\lambda \rangle_t = \int_0^t (w + hf_{2\lambda} - 2h\widehat{m}f_\lambda, \bar{A}_s^\infty) ds.$$

Taking expectation and solving the equation give

$$\mathbb{E}[(f_\lambda, Z_t)] = e^{-ht} \left( e^{\lambda t} \mathbb{E}[(f_\lambda, Z_0)] + \frac{n}{n-\lambda} (e^{nt} - e^{\lambda t}) \mathbb{E}[(1, Z_0)] \right).$$

Suppose that  $\mathbb{E}[Z_0]$  has density  $\mathfrak{z}_0(x)$ , then  $\mathbb{E}[Z_t]$  has density  $\mathfrak{z}_t(x)$  and

$$\mathfrak{z}_t(x) = e^{-ht} \mathfrak{z}_0(x-t) \mathbf{1}_{x>t} + n \mathbb{E}[(1, Z_0)] e^{(n-h)t} e^{-nx} \mathbf{1}_{x \leq t}.$$

The SPDE becomes

$$dZ_t(dx) = (n(1, Z_t) \delta_0(dx) - hZ_t(dx) - (Z_t)'(dx)) dt + d\tilde{M}_t^\infty(dx)$$

with

$$\begin{aligned} \langle \tilde{M}^{f,\infty} \rangle_t &= f^2(0) \frac{w}{n-h} (1, \bar{A}_0^\infty) (e^{(n-h)t} - 1) \\ &\quad + h \int_0^t (f^2, \bar{A}_s^\infty) ds - 2f(0)h\widehat{m} \int_0^t (f, \bar{A}_s^\infty) ds. \end{aligned}$$

In the case where the density exists,

$$\begin{aligned} \langle \tilde{M}^{f,\infty} \rangle_t &= f^2(0) \frac{w}{n-h} (e^{(n-h)t} - 1) \int a_0(x) dx \\ &\quad + h \int_0^t \int f^2(x) a(x, s) dx ds - 2f(0)h\widehat{m} \int_0^t \int f(x) a(x, s) dx ds, \end{aligned}$$

where  $a(x, t) = \begin{cases} a(0, t-x)e^{-hx}, & x \leq t \\ a_0(x-t)e^{-ht}, & x > t \end{cases}$  with  $a(0, t) = n \int a(x, t) dx$  and  $a(x, 0) = a_0(x)$ . In particular,  $\langle \tilde{M}^{1,\infty} \rangle_t = \frac{w+h-2h\widehat{m}}{n-h} (1, \bar{A}_0^\infty) (e^{(n-h)t} - 1)$ .

In fact, we can solve (29) and obtain

$$(f_\lambda, Z_t) = e^{(\lambda-h)t} (f_\lambda, Z_0) + \int_0^t e^{(\lambda-h)(t-s)} (n|Z_s| ds + d\tilde{M}_s^\lambda).$$

With

$$|Z_t| = e^{(n-h)t} |Z_0| + \int_0^t e^{(n-h)(t-s)} d\tilde{M}_s^0,$$

we can write

$$\begin{aligned} (f_\lambda, Z_t) &= e^{(\lambda-h)t} (f_\lambda, Z_0) + \int_0^t e^{(\lambda-h)(t-s)} \left( n(e^{(n-h)s} |Z_0| \right. \\ &\quad \left. + \int_0^s e^{(n-h)(s-r)} d\tilde{M}_r^0) ds + d\tilde{M}_s^\lambda \right) \\ &= e^{(\lambda-h)t} (f_\lambda, Z_0) + \frac{n}{n-\lambda} |Z_0| (e^{(n-h)t} - e^{(\lambda-h)t}) \\ &\quad + ne^{(\lambda-h)t} \int_0^t e^{(n-\lambda)s} \int_0^s e^{-(n-h)r} d\tilde{M}_r^0 ds + \int_0^t e^{(\lambda-h)(t-s)} d\tilde{M}_s^\lambda, \end{aligned}$$

where

$$\langle \tilde{M}^0, \tilde{M}^\lambda \rangle_t = \int_0^t (w - h\widehat{m} + h(1 - \widehat{m})f_\lambda, \bar{A}_s^\infty) ds.$$

## REFERENCES

- [1] R.A. Adams, J.J.F. Fournier (2003). *Sobolev Spaces*, 2nd ed. Elsevier.
- [2] D. Aldous (1978). Stopping times and tightness. *The Annals of Probability* **6**(2), 335–340.
- [3] V. Bansaye, J. Delmas, L. Marsalle, V.C. Tran (2011). Limit theorems for Markov processes indexed by continuous time Galton-Watson trees. *The Annals of Applied Probability* **21**(6), 2263–2314.
- [4] A.-M. Borde-Boussion (1990). Stochastic demographic models: Age of a population. *Stochastic Processes and their Application* **35**, 279–291.
- [5] C. Dellacherie, P.A. Meyer (1978) Probabilities and Potential. North-Holland.
- [6] R. Ferrière, V.C. Tran (2009). Stochastic and deterministic models for age-structured populations with genetically variable traits. *ESAIM: Proceedings, EDP Sciences* **27**, 289–310.
- [7] K. Hamza, P. Jagers, F.C. Klebaner (2013). The age structure of population-dependent general branching processes in environments with a high carrying capacity. *Proceedings of the Steklov Institute of Mathematics* **282**, 90–105.
- [8] T.E. Harris (1963), *The Theory of Branching Processes*. Springer.
- [9] J. Jacod, A.N. Shiryaev (2003). *Limit Theorems for Stochastic Processes*, 2nd ed. Springer.
- [10] P. Jagers (1975). *Branching Processes with Biological Applications*. Wiley. Chichester.
- [11] P. Jagers (1989). General branching processes as Markov fields. *Stochastic Processes and their Applications* **32**, 183–242.
- [12] P. Jagers, F.C. Klebaner (2000). Population-size-dependent and age-dependent branching processes. *Stochastic Processes and Their Applications* **87**, 235–254.
- [13] P. Jagers, F.C. Klebaner (2011). Population-size-dependent, age-structured branching processes linger around their carrying capacity. *Journal of Applied Probability* **48**, 249–260.
- [14] A. Joffe, M. Metivier (1986). Weak convergence of sequences of semimartingales with applications to multitype branching processes. *Advances in Applied Probability* **18**, 20–65.
- [15] H. Kaspi, K. Ramanan (2011). Law of large numbers limits for many-server queues. *The Annals of Applied Probability* **21**(1), 33–114.
- [16] H. Kaspi, K. Ramanan (2013). SPDE limits of many-server queues. *The Annals of Applied Probability* **23**(1), 145–229.
- [17] F.C. Klebaner (1993). Population-dependent branching processes with a threshold. *Stochastic Processes and Their Applications* **46**, 115–127.
- [18] F.C. Klebaner, O. Nerman. (1994) Autoregressive approximation in branching processes with a threshold. *Stochastic Processes and Their Applications* **51**, 1–7.
- [19] S. Meleard (1998). Convergence of the fluctuations for interacting diffusions with jumps associated with Boltzmann equations. *Stochastics and Stochastic Reports* **63**, 195–225.
- [20] S. Meleard, V.C. Tran (2012). Slow and fast scales for superprocess limits of age-structured populations. *Stochastic Processes and Their Applications* **122**, 250–276.
- [21] M. Metivier (1987). Weak convergence of measure valued processes using Sobolev-imbedding techniques. *Lecture Notes in Mathematics* **1236**, 172–183. Springer, Berlin.
- [22] I. Mitoma (1983). On the sample continuity of  $S'$ -processes. *Journal of the Mathematical Society of Japan* **35**(4), 629–636.
- [23] I. Mitoma (1983). Tightness of probability on  $C([0, 1]; S')$  and  $D([0, 1]; S')$ . *The Annals of Probability* **11**(4), 989–999.
- [24] K. Oelschläger (1990). Limit theorem for age-structures populations. *The Annals of Probability* **18**(1), 290–318.
- [25] V.C. Tran (2006). *Modèles particuliers stochastiques pour des problèmes d'évolution adaptative et pour l'approximation de solutions statistiques*. PhD thesis, Université Paris X - Nanterre.
- [26] V.C. Tran (2008). Large population limit and time behaviour of a stochastic particle model describing an age-structured population. *ESAIM: Probability and Statistics* **12**, 345–386.
- [27] V.S. Varadarajan (1958). Weak convergence of measures on separable metric spaces. *Sankhya: The Indian Journal of Statistics*, 19(1/2), pp. 15–22.
- [28] J.B. Walsh (1986). An Introduction to stochastic partial differential equations. In *École d'Été Probabilités de Saint-Flour XIV. Lecture Notes in Mathematics* **1180**, 265–439. Springer, Berlin.

SCHOOL OF MATHEMATICAL SCIENCES, MONASH UNIVERSITY, CLAYTON, VIC. 3058, AUSTRALIA.

*E-mail address:* `kais.hamza@monash.edu`

SCHOOL OF MATHEMATICAL SCIENCES, MONASH UNIVERSITY, CLAYTON, VIC. 3058, AUSTRALIA.

*E-mail address:* `jieyen.fan@monash.edu`

SCHOOL OF MATHEMATICAL SCIENCES, MONASH UNIVERSITY, CLAYTON, VIC. 3058, AUSTRALIA.

*E-mail address:* `kais.hamza@monash.edu`

DEPARTMENT OF MATHEMATICAL SCIENCES, CHALMERS UNIVERSITY OF TECHNOLOGY.

*E-mail address:* `jagers@chalmers.se`

SCHOOL OF MATHEMATICAL SCIENCES, MONASH UNIVERSITY, CLAYTON, VIC. 3058, AUSTRALIA.

*E-mail address:* `fima.klebaner@monash.edu`